

The Dirac operator on collapsing S^1 -bundles

Bernd Ammann

October 1998

Abstract

We study the behavior of the spectrum of the Dirac operator on collapsing S^1 -bundles. Convergent eigenvalues will exist if and only if the spin structure is projectable.

Keywords: Dirac operator, circle bundles, collapse, projectable spin structures

Mathematics Classification: 58G25, 58G30, 53C25

1 Introduction

In this paper we study the spectrum of the Dirac operator on collapsing S^1 -bundles.

There are some nice results about the behavior of the Laplace operator acting on functions on a family of collapsing manifolds. Fukaya [9] proved that if a family $(M_n, \tilde{g}_n)_{n \in \mathbb{N}}$ of Riemannian manifolds with sectional curvature bounded from above and below and bounded diameter converges in the measured Gromov-Hausdorff topology to a Riemannian manifold (N, g) of lower dimension, then the eigenvalues of the Laplace operator acting on functions on (M_n, \tilde{g}_n) converge to those of (N, g) .

Fukaya conjectured that it should be possible to replace the bound on the sectional curvature by a lower bound on the Ricci curvature. So it seems that the connection between Gromov-Hausdorff topology and the spectrum should be much closer than we know until today.

Other papers connecting metric and topological properties to the behavior of the spectrum of the Laplace operator acting on functions are [6], [7] and [14].

For the Laplace operator acting on p -forms with $p \geq 1$ the situation is more complicated. Until now for p -forms there is no analogue to Fukaya's result. But if $\pi : (M, \tilde{g}) \rightarrow (N, g)$ is a Riemannian submersion whose fibers are minimal submanifolds and if the horizontal distribution is integrable, then according to Gilkey and

Park [10, 11] the spectrum of the p -form Laplacian on N is contained in the spectrum of the p -form Laplacian on M . The pullbacks of eigenforms on N are eigenforms on M to the same eigenvalues. So if we have a family of such submersions with fixed (N, g) , then this result trivially implies the convergence of certain eigenvalues.

In a recent paper [12] Gilkey, Leahy and Park generalized some weaker version of this result to arbitrary Riemannian submersions. If an eigenform on N to the eigenvalue λ pulls back to an eigenform on M to the eigenvalue μ , then $\lambda \leq \mu$.

In this paper we will concentrate on another elliptic differential operator, the Dirac operator. We will describe the behavior of the spectrum of the Dirac operator when the fibers of a principal S^1 -bundle collapse. While the Laplacian acting on functions and on forms only depends on the Riemannian metric, the Dirac operator also depends on the spin structure. It will turn out that for some spin structures, the projectable ones, there are convergent eigenvalues whereas for other spin structures all eigenvalues diverge.

The methods of this article generalize the methods of [3] where we only treated the case of geodesic fibers. There are even families that do not have bounded curvature but for which our results imply convergence or non-convergence. Our method is based on a splitting of the Dirac operator into a horizontal Dirac operator, a vertical Dirac operator and a zero order term. An analogous splitting of the Laplace operator has been used in [5].

Unfortunately the generalization to non-geodesic fibers has a drawback: unlike in the geodesic case [3] we do not get any information about the signs of the non-convergent eigenvalues of the Dirac operator.

In the proofs of this paper I omitted some technical calculations and I only dealt with the case when the dimension b of the base manifold is even. More details and the modifications for the odd-dimensional case can be found in my thesis [2]. I thank my supervisor Christian Bär for many interesting discussions and good ideas how to present the collapse result. I also want to thank Bruno Colbois who invited me to Chambéry and Grenoble in November 1997. He told me much about the behavior of the eigenvalues of the Laplacian when the manifolds collapse and much about small eigenvalues of the Laplace operator.

2 Spin structures on S^1 -bundles

We assume that S^1 acts freely and isometrically on a compact, connected, oriented $(b + 1)$ -dimensional Riemannian manifold (M, \tilde{g}) , $b \geq 1$. Then M is the total space of a principal S^1 -bundle over some base space $N^b := M^{b+1}/S^1$. There is a unique metric g on N such that $\pi : (M, \tilde{g}) \rightarrow (N, g)$ is a Riemannian submersion.

The principal S^1 -bundle $M \rightarrow N$ carries a unique connection-1-form

$$i\omega : TM \rightarrow i\mathbb{R},$$

such that $\ker \omega|_m$ is perpendicular to the fibers for any $m \in M$.

The S^1 -action induces a Killing vector field K . The fibers of $M \rightarrow N$ are (totally) geodesic if and only if $\ell := |K|$ is constant in $m \in M$. The length of a fiber is $2\pi\ell$. The metric \tilde{g} on M is completely characterized by ω , ℓ and g .

Now we recall the notion of a spin structure in order to fix the notation. The bundle of positively oriented orthonormal frames $P_{\text{SO}}(N)$ is a principal $\text{SO}(b)$ -bundle. The unique non-trivial double covering of $\text{SO}(b)$ will be denoted $\Theta : \text{Spin}(b) \rightarrow \text{SO}(b)$. A pair $(P_{\text{Spin}}(N), \vartheta)$ will be called *spin structure* if $P_{\text{Spin}}(N)$ is a principal $\text{Spin}(b)$ -bundle over N and if $\vartheta : P_{\text{Spin}}(N) \rightarrow P_{\text{SO}}(N)$ is a Θ -equivariant, fiber preserving map. Two spin structures $(P_{\text{Spin}}^1(N), \vartheta_1)$ and $(P_{\text{Spin}}^2(N), \vartheta_2)$ are isomorphic if there is a $\text{Spin}(b)$ -equivariant, fiber preserving isomorphism $A : P_{\text{Spin}}^1(N) \rightarrow P_{\text{Spin}}^2(N)$ such that $\vartheta_1 = \vartheta_2 \circ A$.

A spin structure exists if and only if the second Stiefel-Whitney class $w_2(TN)$ vanishes. In general the spin structure is not unique. Having chosen a spin structure we can define a spinor bundle on N , Clifford multiplication and a Dirac operator. The spectrum of the Dirac operator depends on the choice of spin structure.

The definitions of spin structure, Dirac operator, ... on M are clearly analogous. The S^1 -action on M induces an S^1 -action on $P_{\text{SO}}(M)$. A spin structure $\vartheta : P_{\text{Spin}}(M) \rightarrow P_{\text{SO}}(M)$ is said to be *projectable* if this S^1 -action lifts continuously to $P_{\text{Spin}}(M)$.

Examples.

- (1) We view the $(b + 1)$ -dimensional torus T^{b+1} as the total space of the circle bundle $T^{b+1} \rightarrow T^b$. The torus T^{b+1} carries 2^{b+1} spin structures, half of them are projectable, half of them are non-projectable. Recently the special case $T^2 \rightarrow T^1 = S^1$ with the trivial spin structure and non-geodesic fibers has been intensively studied in [1].
- (2) We view S^{2l+1} as the total space of the Hopf fibration $S^{2l+1} \rightarrow \mathbb{C}P^l$. The (unique) spin structure on S^{2l+1} is projectable if and only if $\mathbb{C}P^l$ is spin and therefore if and only if l is odd.

There is a natural isomorphism from projectable spin structures on M to spin structures on N . However M may admit a non-projectable spin structure, even if there are no spin structures on N . The Hopf fibration $S^{2l+1} \rightarrow \mathbb{C}P^l$ with l even is an example for this phenomenon.

In this paper we will not only look at one single S^1 -fibered space M , but at a family M_n of such spaces over a base space N that does not depend on n . For each $n \in \mathbb{N}$ the manifold M_n carries a Riemannian metric \tilde{g}_n , such that $\pi_n : (M_n, \tilde{g}_n) \rightarrow (N, g)$ is an S^1 -bundle and a Riemannian submersion. The quantities ℓ_n and ω_n are defined as they are defined for M . Roughly speaking, we will analyze the behavior of the spectrum of the Dirac operator if ℓ_n tends to 0, whereas ω_n stays small in a suitable sense. This situation will be called *collapse*.

Note that N and g do not depend on n . On the other hand M_n may even change its topological type for different values of n .

3 Projectable spin structures

Let (N, g) be a Riemannian manifold carrying a spin structure that shall be fixed throughout this section. The eigenvalues of the Dirac operator D on N will be denoted by $(\mu_j)_{j \in \mathbb{N}}$. Moreover let $(M_n, \tilde{g}_n)_{n \in \mathbb{N}}$ be a family of Riemannian manifolds such that S^1 acts freely and isometrically on each M_n , and let $(M_n, \tilde{g}_n) \rightarrow (N = M_n/S^1, g)$ be Riemannian submersions.

The spin structure on M_n shall be the unique projectable spin structure corresponding to the spin structure on N . As above let $2\pi\ell_n$ be the length of the fibers and $i\omega_n$ the connection-1-form on $M_n \rightarrow N$. Furthermore we assume the collapsing condition

$$\begin{aligned} \|\ell_n \cdot d\omega_n\|_\infty &\rightarrow 0 \text{ and } \|\ell_n\|_\infty \rightarrow 0 \text{ for } n \rightarrow \infty \\ \alpha &:= \limsup_{n \rightarrow \infty} \|\text{grad} \ell_n\|_\infty < 1 \end{aligned} \tag{1}$$

THEOREM 3.1. *The eigenvalues $(\lambda_{j,k}(n))_{j \in \mathbb{N}, k \in \mathbb{Z}}$ of the Dirac operator \tilde{D}^n on M_n can be numbered in such a way that:*

- (1) *For all $\varepsilon > 0$ there is a $n_0 \in \mathbb{N}$, such that we have for any $n \geq n_0$ and $j \in \mathbb{N}$, $k \in \mathbb{Z} - \{0\}$*

$$\|\ell_n\|_\infty^2 \lambda_{j,k}(n)^2 \geq |k| (|k| - \alpha) - \varepsilon.$$

In particular we get $\lambda_{j,k}(n)^2 \rightarrow \infty$ for $n \rightarrow \infty$.

Furthermore, if M_n and ω_n do not depend on n , then we also have for $j \in \mathbb{N}$, $k \in \mathbb{Z} - \{0\}$.

$$\limsup_{n \rightarrow \infty} \left(\min_{p \in N} \ell_n(p) \right)^2 \lambda_{j,k}(n)^2 \leq |k| (|k| + \alpha).$$

This upper bound of $\lambda_{j,k}(n)^2$ is not uniform in j and k .

(2) If $b = \dim N$ is even, then we get for $n \rightarrow \infty$

$$\lambda_{j,0}(n) \rightarrow \mu_j.$$

However for $b = \dim N$ odd we obtain

$$\begin{aligned} \lambda_{2j-1,0}(n) &\rightarrow \mu_j \\ \lambda_{2j,0}(n) &\rightarrow -\mu_j \end{aligned}$$

In both cases the convergence of the eigenvalues $\lambda_{j,0}(n)$ is uniform in j .

Examples.

(1) The square of the Dirac operator on the flat torus $\mathbb{R}/a\mathbb{Z} \times \mathbb{R}/\ell\mathbb{Z}$ with the trivial spin structure has the eigenvalues

$$\lambda_{j_1, j_2, k}^2 = 4\pi^2 \left(\frac{j_1^2}{a^2} + \frac{k^2}{\ell^2} \right), \quad j_1 \in \mathbb{Z}, j_2 \in \{0, 1\}, k \in \mathbb{Z}.$$

(2) We take the standard metric on $\mathbb{C}P^l$, the ω coming from the Hopf fibration $S^{2l+1} \rightarrow \mathbb{C}P^l$ and a sequence of constant functions $(\ell_n)_{n \in \mathbb{N}}$, $\ell_n > 0$, $\ell_n \rightarrow 0$. The metrics on S^{2l+1} characterized by these quantities are called *Berger metrics* g_{ℓ_n} . If l is odd, some eigenvalues diverge whereas other eigenvalues converge to the eigenvalues of $\mathbb{C}P^l$. The spectrum of the spheres with Berger metrics has been explicitly calculated in [13, 4]. C. Bär used the collapse results to calculate the spectrum of the Dirac operator on $\mathbb{C}P^l$ [3].

Idea of proof. In order to prove Theorem 3.1 we write the Dirac operator \widetilde{D}^n as a sum of a “vertical Dirac operator”, a “horizontal Dirac operator” and a zero order term.

For the definition of the vertical Dirac operator we need the Lie derivative of spinors in the direction of the Killing field. The S^1 -action on $P_{\text{Spin}}M_n$ induces an S^1 -action on $\Sigma M_n = P_{\text{Spin}}(M_n) \times_{\text{Spin}(b+1)} \Sigma_{b+1}$. The latter action will be denoted by κ . A spinor with base point $m \in M$ will be mapped by $\kappa(e^{is})$ to a spinor with base point $m \cdot e^{is}$. We define the *Lie derivative* of a smooth spinor field Ψ in the direction of the Killing field K as

$$\mathcal{L}_K(\Psi)(m) = \left. \frac{d}{ds} \right|_{s=0} \kappa(e^{-is})(\Psi(m \cdot e^{is})).$$

As \mathcal{L}_K is the differential of a representation of the Lie group S^1 on $L^2(\Sigma M_n)$, we get the decomposition

$$L^2(\Sigma M_n) = \bigoplus_{k \in \mathbb{Z}} V_{k,n}$$

into the eigenspaces $V_{k,n}$ of the operator \mathcal{L}_K to the eigenvalue ik , $k \in \mathbb{Z}$. The S^1 -action commutes with the Dirac operator on M_n , and therefore this decomposition is respected by the Dirac operator.

We calculate the difference between the covariant derivative and the Lie derivative in the direction of K . For any smooth section $\tilde{\Psi}$ of ΣM_n we get

$$\nabla_K \tilde{\Psi} - \mathcal{L}_K \tilde{\Psi} = \frac{\ell_n^2}{4} \gamma(d\omega_n) \tilde{\Psi} - \frac{1}{2} \gamma(K/\ell_n) \gamma(\text{grad} \ell_n) \tilde{\Psi}. \quad (2)$$

Here $\gamma(V)$ resp. $\gamma(\beta)$ denotes Clifford multiplication of a spinor by the vector V and the 2-form β resp.

What we got until now is some kind of Fourier decomposition along the fibers. In the following it will turn out that for any k there is a natural isomorphism $Q_{k,n}$ from the summand $V_{k,n}$ to the vector space of sections of a twisted spinor bundle on the quotient space N .

To the principal S^1 -bundle $\pi_n : M_n \rightarrow N$ we associate the complex vector bundle $L_n := M_n \times_{S^1} \mathbb{C}$ with a connection given by $i\omega_n$. The horizontal lift of a vector (field) X on N to a vector (field) on M_n will be denoted by \tilde{X} .

For b even, the isomorphism $Q_{k,n}$ is given by the following technical lemma that is proven in [2, Lemma-Definition 7.2.3].

LEMMA 3.2. *Let $b = \dim N$ be even. Then for any n there is an isometry of Hilbert spaces*

$$Q_{k,n} : L^2(\Sigma N \otimes L_n^{-k}) \rightarrow V_{k,n}$$

such that the horizontal covariant derivative is given by

$$\nabla_{\tilde{X}} Q_{k,n}(\Psi) = Q_{k,n}(\nabla_X \Psi) + \frac{\ell_n}{4} \gamma(K/\ell_n) \gamma(\tilde{W}_X) Q_{k,n}(\Psi) - \frac{X(\ell_n)}{2\ell_n} Q_{k,n}(\Psi),$$

where W_X is the vector field on N satisfying $d\omega_n(\tilde{X}, \cdot) = \langle \tilde{W}_X, \cdot \rangle$. Clifford multiplication is preserved, i. e.

$$Q_{k,n}(\gamma(X)\Psi) = \gamma(\tilde{X})Q_{k,n}(\Psi).$$

Now we are ready to prove the theorem for b even. We define the *horizontal Dirac operator* as the unique closed linear operator $D_h^n : L^2(\Sigma M_n) \rightarrow L^2(\Sigma M_n)$ on each $V_{k,n}$ given by

$$D_h^n := Q_{k,n} \circ D^n \circ Q_{k,n}^{-1}$$

where D^n is the twisted Dirac operator on $\Sigma N \otimes L_n^{-k}$.

We define the *vertical Dirac operator*

$$D_v^n := \gamma(K/\ell_n) \mathcal{L}_K$$

and the zero order term

$$Z_n := -(1/4) \gamma(K/\ell_n) \gamma(d\omega_n).$$

Using Formula (2) and Lemma 3.2 we can express the Dirac operator as a sum:

$$\widetilde{D}^n = \frac{1}{\ell_n} D_v^n + D_h^n + \ell_n Z_n.$$

Since D_h^n , $\gamma(K/\ell_n)$ and Z_n commute with the S^1 -action they also commute with \mathcal{L}_K . Therefore each summand of the above decomposition of the Dirac operator maps each $V_{k,n}$ into itself. Furthermore the spectrum of $D_h^n|_{V_{0,n}}$ is just the spectrum of the Dirac operator D acting on sections of ΣN . So the eigenvalues of $\widetilde{D}^n|_{V_{0,n}}$ converge to the eigenvalues of D . This fact immediately implies (2) of the theorem for b even.

An elementary calculation shows that D_h^n anticommutes with $\gamma(K/\ell_n)$ and therefore it anticommutes with $D_v^n := \gamma(K/\ell_n)\mathcal{L}_K$.

From this anticommutativity we get the formula

$$D_h^n \left(\frac{1}{\ell_n} D_v^n \right) + \left(\frac{1}{\ell_n} D_v^n \right) D_h^n = \gamma(\text{grad}(1/\ell_n)) D_v^n.$$

We take the square of $A^n := (1/\ell_n)D_v^n + D_h^n$ and get

$$(A^n)^2 = \frac{1}{\ell_n^2} (D_v^n)^2 + (D_h^n)^2 - \frac{\gamma(\text{grad } \ell_n)}{\ell_n^2} D_v^n.$$

Now for $\alpha_n := \|\text{grad } \ell_n\|_\infty$ and for $\tilde{\Psi} \in V_{k,n}$, $k \neq 0$ we have

$$\begin{aligned} ((A^n)^2 \tilde{\Psi}, \tilde{\Psi})_{M_n} &= \left(\left(\frac{k^2}{\ell_n^2} + (D_h^n)^2 - ik \frac{\gamma(\text{grad } \ell_n) \gamma(K/\ell_n)}{\ell_n^2} \right) \tilde{\Psi}, \tilde{\Psi} \right)_{M_n} \\ &\geq \frac{|k|(|k| - \alpha_n)}{\|\ell_n\|_\infty^2} (\tilde{\Psi}, \tilde{\Psi})_{M_n}. \end{aligned}$$

The eigenvalues of $(A^n)^2|_{V_{k,n}}$ are therefore larger than or equal to

$$\left(|k| (|k| - \alpha_n) \right) / \|\ell_n\|_\infty^2.$$

As $\ell_n \cdot d\omega_n$ is a bounded operator, whose norm vanishes for $n \rightarrow \infty$, the norm of the zero order term tends to zero. So we get the first part of (1) of the theorem.

Now let M_n and ω_n be independent from n . Then also L_n and D^n are independent from n . Analogously we obtain

$$\begin{aligned} \left((A^n)^2 Q_{k,n}(\Psi), Q_{k,n}(\Psi) \right)_{M_n} &\leq \left((D^n)^2 \Psi, \Psi \right)_N \\ &+ \frac{|k|(|k| + \alpha_n)}{\left(\min_{p \in N} \ell_n(p) \right)^2} \left(Q_{k,n}(\Psi), Q_{k,n}(\Psi) \right)_{M_n} \end{aligned}$$

From this we get the second part of (1) for b even.

The proof for b odd runs quite analogously, but we have to do some modifications as in this case the rank of the spinor bundle over M_n is two times the rank of the spinor bundle over N . For details about this case we refer to [3] or [2]. \square

Remark. Suppose we have a complex vector bundle $E \rightarrow N$ with a metric connection ∇^E . Now we replace the Dirac operator D by the twisted Dirac operator D^E acting on sections of $\Sigma N \otimes E \rightarrow N$, the Dirac operator \widetilde{D}^n will be replaced by the twisted Dirac operator $\widetilde{D}^{n,E}$ acting on sections of $\Sigma M_n \otimes \pi_n^* E \rightarrow M_n$. Then Theorems 3.1, 4.1 and 4.2 are still valid. The proof of this generalized version is essentially the same as the proof above.

4 Non-projectable spin structures

Now we turn to the case of non-projectable spin structures. In this case we get a similar result, but with a stronger restriction on the gradient of the fiber length. The variable k from the last section does no longer take integer values, but values in $\mathbb{Z} + (1/2)$. Therefore all eigenvalues will diverge.

We define N , g , M_n , \tilde{g}_n , ω_n and ℓ_n as above. However in contrast to the last section we do not assume that N carries any spin structure. Instead we suppose that each M_n carries a non-projectable spin structure.

The collapsing condition in this section is

$$\begin{aligned} \|\ell_n \cdot d\omega_n\|_\infty &\rightarrow 0 \text{ and } \|\ell_n\|_\infty \rightarrow 0 \text{ for } n \rightarrow \infty \\ \alpha &:= \limsup_{n \rightarrow \infty} \|\text{grad } \ell_n\|_\infty < 1/2. \end{aligned} \tag{3}$$

THEOREM 4.1. *All eigenvalues of the Dirac operator \widetilde{D}^n on M_n diverge.*

Moreover, the eigenvalues $(\lambda_{j,k}(n))_{j \in \mathbb{N}, k \in \mathbb{Z} + (1/2)}$ can be numbered in such a way that

(a) For any $\varepsilon > 0$ there is $n_0 \in \mathbb{N}$, such that for any $n \geq n_0$ and $j \in \mathbb{N}$, $k \in \mathbb{Z} + (1/2)$ we get

$$\|\ell_n\|_\infty^2 \lambda_{j,k}(n)^2 \geq |k| (|k| - \alpha) - \varepsilon.$$

(b) If M_n and ω_n are independent from n , then we additionally have for any $j \in \mathbb{N}$, $k \in \mathbb{Z} + (1/2)$

$$\limsup_{n \rightarrow \infty} \left[\left(\min_{p \in N} \ell_n(p) \right)^2 \lambda_{j,k}(n)^2 \right] \leq |k| (|k| + \alpha).$$

This upper bound is not uniform in j and k .

Example. Suppose we have a family of collapsing Berger metrics on S^{2l+1} as in the previous example, but with l even. Then all eigenvalues diverge. The same holds for tori with non-projectable spin structures.

Idea of proof. The proof is a variation of the proof of Theorem 3.1. We will restrict to the case b even.

Let $\tilde{\vartheta}_n : P_{\text{Spin}}(M_n) \rightarrow P_{\text{SO}}(M_n)$ be a non-projectable spin structure on M_n . In this case N may or may not be spin. We define $P_{\text{SO}(b)}(M_n)$ to be the set of all frames in $P_{\text{SO}}(M_n)$ having K/ℓ_n as first vector. Then $P_{\text{SO}(b)}(M_n)$ is a principal $\text{SO}(b)$ -bundle over M_n . Moreover $P := \tilde{\vartheta}_n^{-1}(P_{\text{SO}(b)}(M_n))$ is a principal $\text{Spin}(b)$ -bundle over M_n .

The action of $S^1 \cong (\mathbb{R}/2\pi\mathbb{Z})$ does not lift to P , but the double covering of S^1 , i.e. $S^1 \cong (\mathbb{R}/4\pi\mathbb{Z})$, does act on P . We define $\text{Spin}^{\mathbb{C}}(b)$ to be $\text{Spin}(b) \times_{\mathbb{Z}_2} (\mathbb{R}/4\pi\mathbb{Z})$ where $-1 \in \mathbb{Z}_2$ identifies $(-A, c)$ with $(A, c + 2\pi)$. The complex standard representation Σ_b of $\text{Spin}(b)$ is also a representation for $\text{Spin}^{\mathbb{C}}(b)$ where $s + 4\pi\mathbb{Z} \in \mathbb{R}/4\pi\mathbb{Z}$ operates as $\exp(is/2)$. The actions of $\text{Spin}(b)$ and $\mathbb{R}/4\pi\mathbb{Z}$ on P induce a free action of $\text{Spin}^{\mathbb{C}}(b)$ on P and we can view P as a principal $\text{Spin}^{\mathbb{C}}(b)$ -bundle over N . Then we can form the bundle $P \times_{\text{Spin}^{\mathbb{C}}(b)} \Sigma_b$.

If N is spin, this bundle is just $\Sigma N \otimes L_n^{\frac{1}{2}}$. If N is not spin, then neither ΣN nor $L_n^{\frac{1}{2}}$ exist but $\Sigma N \otimes L_n^{\frac{1}{2}}$ does exist.

Again we get a splitting

$$L^2(\Sigma M_n) = \bigoplus_{k \in \mathbb{Z} + \frac{1}{2}} V_{k,n}$$

into eigenspaces $V_{k,n}$ for \mathcal{L}_K to the eigenvalue ik . The rest of the proof of Theorem 4.1 is the same as the one for the case $k \neq 0$ in Theorem 3.1. \square

The formulas in the proof also show the following lower bound for eigenvalues of D^2 . We define the *Clifford norm* $\|\eta\|_{\text{Cl}}$ of a 2-form η on M to be the L^2 -operator norm of $\gamma(\eta) \in \text{End}(\Gamma \Sigma M)$.

THEOREM 4.2. *Let (M, \tilde{g}) be a Riemannian manifold on which S^1 acts freely and isometrically. The quotient $N := M/S^1$ shall carry the unique metric g for which $M \rightarrow N$ is a Riemannian submersion. We assume that M carries a non-projectable spin structure.*

If $\alpha := \|\text{grad } \ell\|_\infty < 1/2$, then all eigenvalues $(\lambda_i)_{i \in \mathbb{Z}}$ of the Dirac operator \tilde{D} on M satisfy

$$|\lambda_i| \geq \frac{\sqrt{1 - 2\|\text{grad } \ell\|_\infty}}{2\|\ell\|_\infty} - \frac{\|\ell d\omega\|_{\text{Cl}}}{4}.$$

Remark. This lower bound for the smallest eigenvalue of the Dirac operator depends on the spin structure.

The existence of such a bound is not surprising: on many standard spaces the spectrum of the Dirac operator is explicitly known, among many others flat tori [8] and compact quotients of 3-dimensional Heisenberg groups with a left invariant metric [3]. For these examples it turns out that the spectrum strongly depends on the spin structure, in particular the smallest eigenvalue also depends on it. So for metrics close enough to these standard spaces there should be a lower bound for the smallest eigenvalue of the Dirac operator depending on the spin structure.

Until recently such a bound was not known. Theorem 4.2 provides such a bound for the case of a circle bundle with non-projectable spin structure and sufficiently short fibers. If $\|\text{grad } \ell\|_\infty$ and $\|\ell\|_\infty$ are sufficiently small Theorem 4.2 gives a better bound than any other estimate published before. For more details and other recent estimates depending on the spin structure I refer to my thesis [2].

References

- [1] I. Agricola, B. Ammann, T. Friedrich, *A comparison of the eigenvalues of the Dirac and Laplace operator on the two-dimensional torus*, Preprint 1998
- [2] B. Ammann, *Spin-Strukturen und das Spektrum des Dirac-Operators*, Dissertation, Universität Freiburg, ISBN 3-8265-4282-7, Shaker Verlag Aachen 1998
- [3] B. Ammann, C. Bär, *The Dirac Operator on Nilmanifolds and Collapsing Circle Bundles*, Ann. Global Anal. Geom. **16**, 221–253 (1998)
- [4] C. Bär, *Metrics with harmonic spinors*, GAFA **6**, 899–942 (1996)
- [5] L. Bérard Bergery, J.-P. Bourguignon, *Riemannian submersions with totally geodesic fibers*, Illinois J. Math. **26**, 181–200 (1982)

- [6] B. Colbois, G. Courtois, *Convergence de variétés et convergence du spectre du Laplacien*, Ann. Sc. Ec. Norm. Sup. 4^e série **24**, 507–518 (1991)
- [7] B. Colbois, J. Dodziuk, *Riemannian metrics with large λ_1* , Proc. Amer. Math. Soc. **122** no. 3, 905–906 (1994)
- [8] T. Friedrich, *Zur Abhängigkeit des Dirac-Operators von der Spin-Struktur*, Colloq. Math. **48**, 57–62 (1984)
- [9] K. Fukaya, *Collapsing of Riemannian manifolds and eigenvalues of Laplace operator*, Invent. Math. **87**, 517–547 (1987)
- [10] P. B. Gilkey, J. H. Park, *Eigenvalues of the Laplacian and Riemannian submersions*, Yokohama Math. J. **43** no. 1, 7–11 (1995)
- [11] P. B. Gilkey, J. H. Park, *Riemannian submersions which preserve the eigenforms of the Laplacian*, Ill. J. Math. **40**, 194–201 (1996)
- [12] P. B. Gilkey, J. V. Leahy, J. H. Park, *Eigenvalues of the form valued Laplacian for Riemannian submersions*, Proc. Amer. Math. Soc. **126** no. 6, 1845–1850 (1998)
- [13] N. Hitchin, *Harmonic spinors*, Adv. Math. **14**, 1-55 (1974)
- [14] J.-Y. Wu, *The topological spectrum of a smooth closed manifold*, Ind. Univ. Math. J. **44**, no. 2, 511–534 (1995)