

# SURGERY AND THE SPINORIAL $\tau$ -INVARIANT

BERND AMMANN, MATTIAS DAHL, AND EMMANUEL HUMBERT

ABSTRACT. We associate to a compact spin manifold  $M$  a real-valued invariant  $\tau(M)$  by taking the supremum over all conformal classes of the infimum inside each conformal class of the first positive Dirac eigenvalue, when the metrics are normalized to unit volume. This invariant is a spinorial analogue of Schoen's  $\sigma$ -constant, also known as the smooth Yamabe invariant.

We prove that if  $N$  is obtained from  $M$  by surgery of codimension at least 2 then  $\tau(N) \geq \min\{\tau(M), \Lambda_n\}$ , where  $\Lambda_n$  is a positive constant depending only on  $n = \dim M$ . Various topological conclusions can be drawn, in particular that  $\tau$  is a spin-bordism invariant below  $\Lambda_n$ . Also, below  $\Lambda_n$  the values of  $\tau$  cannot accumulate from above when varied over all manifolds of dimension  $n$ .

## CONTENTS

1. Introduction	2
1.1. Spin manifolds and Dirac operators	2
1.2. The $\tau$ -invariant	2
1.3. The $\sigma$ -constant	3
1.4. Geometric constants	3
1.5. Joining manifolds	4
1.6. Surgery and bordism	4
1.7. Variants of the results	6
2. Preliminaries	6
2.1. Notation for balls and neighbourhoods	6
2.2. Joining manifolds along submanifolds	7
2.3. Comparing spinors for different metrics	8
2.4. Regularity results	9
2.5. The associated variational problem	9
3. Preparations for proofs	10
3.1. Removal of singularities	10
3.2. Limit spaces and limit solutions	11
3.3. Dirac spectral bounds on products with spheres	13
3.4. Approximation by local product metrics	13
4. Proofs	17
4.1. Proof of Theorem 1.2	17
4.2. Proof of Theorem 1.1	28

---

*Date:* August 26, 2008.

*2000 Mathematics Subject Classification.* 53C27 (Primary) 55N22, 57R65 (Secondary).

*Key words and phrases.* Dirac operator, eigenvalue, surgery.

We thank Victor Nistor for helpful discussions about Sobolev inequalities on non-compact manifolds. Also Mattias Dahl thanks the Institut Élie Cartan, Nancy, for its kind hospitality and support during visits when this work was initiated.

## 1. INTRODUCTION

**1.1. Spin manifolds and Dirac operators.** Let  $M$  be a compact  $n$ -dimensional spin manifold without boundary. We will always consider spin manifolds as equipped with an orientation and a spin structure. The existence of these structures is equivalent to the vanishing of the first and second Stiefel-Whitney class.

As explained in [19, 10, 14] one associates the spinor bundle  $\Sigma_\rho^g M$  to the spin structure, together with a Riemannian metric  $g$  on  $M$  and a complex irreducible representation  $\rho$  of the Clifford algebra over  $\mathbb{R}^n$ . The Dirac operator  $D_\rho^g$  is a self-adjoint elliptic first order differential operator acting on smooth sections of the spinor bundle  $\Sigma^g M$ . It has a spectrum consisting only of real eigenvalues of finite multiplicity. The spectrum depends on the choice of spin structure, on the metric  $g$ , and a priori on the representation  $\rho$ . In even dimensions  $n$ , the representation  $\rho$  is unique. In odd dimensions there are two choices  $\rho^+$  and  $\rho^-$ . Exchanging the representation results in reversing the spectrum, that is if  $\lambda$  is an eigenvalue of  $D_{\rho^+}^g$  then  $-\lambda$  is an eigenvalue of  $D_{\rho^-}^g$  with the same multiplicity, and vice versa. This has no effect if  $n \equiv 1 \pmod{4}$  since the real/quaternionic structure on  $\Sigma_\rho^g M$  anti-commutes with the Dirac operator and the spectrum therefore is symmetric, see [10, Section 1.7]. However, in dimensions  $n \equiv 3 \pmod{4}$  the choice of  $\rho$  matters. In this case we choose the representation such that Clifford multiplication of  $e_1 \cdot e_2 \cdots e_n$  acts as the identity, where  $e_1, \dots, e_n$  denotes the standard basis of  $\mathbb{R}^n$ . We thus can and will suppress  $\rho$  in the notation.

**1.2. The  $\tau$ -invariant.** We denote by  $\lambda_1^+(D^{\tilde{g}})$  the first non-negative eigenvalue of  $D^{\tilde{g}}$ . For a metric  $g$  on  $M$  we define

$$\lambda_{\min}^+(M, g) := \inf \lambda_1^+(D^{\tilde{g}}) \text{Vol}(M, \tilde{g})^{1/n},$$

where the infimum is taken over all metrics  $\tilde{g}$  conformal to  $g$ . Further we define

$$\tau^+(M) := \sup \lambda_{\min}^+(M, g),$$

where the supremum is taken over all metrics  $g$  on  $M$ . This yields an invariant of the spin manifold  $M$ . Observe that we do not require  $M$  to be connected.

We begin by noting some simple properties of the invariant  $\tau^+$ . Let  $(S^n, \sigma^n)$  denote the unit sphere with its standard metric. We have

$$\lambda_{\min}^+(S^n, \sigma^n) = \frac{n}{2} \omega_n^{1/n},$$

where  $\omega_n$  is the volume of  $(S^n, \sigma^n)$ . Moreover it is shown in [2, 6] that

$$\lambda_{\min}^+(M, g) \leq \lambda_{\min}^+(S^n, \sigma^n)$$

for any compact Riemannian spin manifold  $(M, g)$ . Together with Inequality (1) below we get

$$\tau^+(S^n) = \lambda_{\min}^+(S^n, \sigma^n) = \frac{n}{2} \omega_n^{1/n},$$

so for all compact spin manifolds  $M$  we have

$$\tau^+(M) \leq \tau^+(S^n).$$

If the kernel of  $D^g$  is non-trivial, then obviously  $\lambda_{\min}^+(M, g) = 0$ . Conversely, it was shown in [2] that if the kernel of  $D^g$  is trivial, that is if  $D^g$  is invertible, then  $\lambda_{\min}^+(M, g) > 0$ . It follows that  $\tau^+(M) > 0$  if and only if there is a metric  $g$  on  $M$  for which the Dirac operator  $D^g$  is invertible. It is a further fact that  $\tau^+(M) = 0$  precisely when  $\alpha(M) \neq 0$ , where  $\alpha(M)$  is the alpha-invariant which equals the index of the Dirac operator for any metric on  $M$ , see [4].

For compact Riemannian spin manifolds  $(M_1, g_1)$  and  $(M_2, g_2)$  we denote by  $M_1 \amalg M_2$  the disjoint union of  $M_1$  and  $M_2$  with the natural metric  $g_1 \amalg g_2$ . It is not difficult to see that

$$\lambda_{\min}^+(M_1 \amalg M_2, g_1 \amalg g_2) = \min\{\lambda_{\min}^+(M_1, g_1), \lambda_{\min}^+(M_2, g_2)\}.$$

This implies

$$\tau^+(M_1 \amalg M_2) = \min\{\tau^+(M_1), \tau^+(M_2)\}.$$

We denote by  $-M$  the manifold  $M$  equipped with the opposite orientation. The Dirac operator changes sign when the orientation of the manifold is reversed. If  $M$  has dimension  $\not\equiv 3 \pmod{4}$  this does not change the first positive eigenvalue of  $D$  since the spectrum is symmetric, so we then have  $\lambda_{\min}^+(-M, g) = \lambda_{\min}^+(M, g)$  and  $\tau^+(-M) = \tau^+(M)$ . For manifolds  $M$  of dimension  $\equiv 3 \pmod{4}$  we define  $\lambda_{\min}^-(M, g)$  and  $\tau^-(M)$  similar to  $\lambda_{\min}^+(M, g)$  and  $\tau^+(M)$  by replacing  $\lambda_1^+$  by the absolute value of the first non-positive eigenvalue. We then have  $\lambda_{\min}^+(-M, g) = \lambda_{\min}^-(M, g)$  and  $\tau^+(-M) = \tau^-(M)$ .

**1.3. The  $\sigma$ -constant.** The  $\tau$ -invariant is a spinorial analogue of the  $\sigma$ -constant [17, 21] which is defined for a compact manifold  $M$  by

$$\sigma(M) := \sup \inf \frac{\int \text{Scal}^{\tilde{g}} dv^{\tilde{g}}}{\text{Vol}(M, \tilde{g})^{\frac{n-2}{n}}},$$

where the infimum runs over all metrics  $\tilde{g}$  in a conformal class and the supremum runs over all conformal classes.  $\sigma(M)$  is also known as the smooth Yamabe invariant of  $M$ . When  $\sigma(M)$  is positive it can be computed in a way analogous to  $\tau^+(M)$  using the lowest eigenvalue of the conformal Laplacian  $L^g = 4\frac{n-1}{n-2}\Delta^g + \text{Scal}^g$  instead of  $\lambda_1^+(D^g)$ . Hijazi's inequality [12, 13] gives a comparison of the two invariants,

$$\tau^\pm(M)^2 \geq \frac{n}{4(n-1)}\sigma(M). \quad (1)$$

For  $M = S^n$  equality is attained in (1). Upper bounds for  $\tau^\pm(M)$  may help to determine the  $\sigma$ -constant.

Surgery formulas for the  $\sigma$ -constant analogous to those obtained in this paper have been proved in [5].

**1.4. Geometric constants.** We are going to prove a surgery formula for the invariant  $\tau^+$ . This formula involves geometric constants  $\Lambda_{n,k}$  which we now define.

For a complete spin manifold  $(V, g)$  we set

$$\widetilde{\lambda_{\min}^+}(V, g) := \inf \lambda \in [0, \infty],$$

where the infimum is taken over all  $\lambda \in (0, \infty)$  for which there is a non-zero spinor field  $\varphi \in L^\infty(V) \cap L^2(V) \cap C_{\text{loc}}^1(V)$  such that  $\|\varphi\|_{L^{\frac{2n}{n-1}}(V)} \leq 1$ , and

$$D^g \varphi = \lambda |\varphi|^{\frac{2}{n-1}} \varphi. \quad (2)$$

If there are no such solutions of (2) on  $V$  then  $\widetilde{\lambda}_{\min}^+(V, g) = \infty$ .

For a positive integer  $k$  we let  $\xi^k$  denote the Euclidean metric on  $\mathbb{R}^k$ . For  $c \in \mathbb{R}$  we denote by  $\eta_c^{k+1} := e^{2ct}\xi^k + dt^2$  the hyperbolic metric of sectional curvature  $-c^2$  on  $\mathbb{R}^{k+1}$ . As above  $\sigma^{n-k-1}$  denotes the metric of sectional curvature 1 on  $S^{n-k-1}$ . We define the product metric

$$G_c := \eta_c^{k+1} + \sigma^{n-k-1}$$

on  $\mathbb{R}^{k+1} \times S^{n-k-1}$ , and we define our geometric invariants as

$$\Lambda_{n,k} := \inf_{c \in [-1,1]} \widetilde{\lambda}_{\min}^+(\mathbb{R}^{k+1} \times S^{n-k-1}, G_c),$$

and

$$\Lambda_n := \min_{0 \leq k \leq n-2} \Lambda_{n,k}.$$

Note that the infimum could as well be taken over  $c \in [0, 1]$  since  $G_c$  and  $G_{-c}$  are isometric. It is easy to see that  $\Lambda_{n,0} = \lambda_{\min}^+(S^n, \sigma^n)$ . For  $k > 0$  we are not able to compute these constants, but at least we can show that they are positive.

**Theorem 1.1.** *For  $0 \leq k \leq n - 2$  we have  $\Lambda_{n,k} > 0$ .*

**1.5. Joining manifolds.** We are going to study the behaviour of  $\tau^+$  when two compact Riemannian spin manifolds are joined along a common submanifold. Let  $M_1$  and  $M_2$  be spin manifolds of dimension  $n$  and let  $N$  be obtained by joining  $M_1$  and  $M_2$  along a common submanifold as described in Section 2.2. The manifold  $N$  is spin and from the construction there is a natural choice of spin structure on  $N$ . The following results make it possible to compare  $\tau^+(M_1 \amalg M_2)$  and  $\tau^+(N)$ .

**Theorem 1.2.** *Let  $(M_1, g_1)$  and  $(M_2, g_2)$  be compact Riemannian spin manifolds of dimension  $n$  for which both  $D^{g_1}$  and  $D^{g_2}$  have trivial kernel. Let  $W$  be a compact spin manifold of dimension  $k$  embedded into  $M_1$  and  $M_2$  with trivializations of the corresponding normal bundles given. Assume that  $0 \leq k \leq n - 2$ , and let  $N$  be obtained by joining  $M_1$  and  $M_2$  along  $W$ . Then there is a family of metrics  $g_\theta$ ,  $\theta \in (0, \theta_0)$  on  $N$  satisfying*

$$\begin{aligned} \min\{\lambda_{\min}^+(M_1 \amalg M_2, g_1 \amalg g_2), \Lambda_{n,k}\} &\leq \liminf_{\theta \rightarrow 0} \lambda_{\min}^+(N, g_\theta) \\ &\leq \limsup_{\theta \rightarrow 0} \lambda_{\min}^+(N, g_\theta) \\ &\leq \lambda_{\min}^+(M_1 \amalg M_2, g_1 \amalg g_2). \end{aligned}$$

Taking the supremum over all metrics on  $M_1 \amalg M_2$  the first inequality gives us the following corollary.

**Corollary 1.3.** *In the situation of Theorem 1.2 we have*

$$\tau^+(N) \geq \min\{\tau^+(M_1 \amalg M_2), \Lambda_{n,k}\} \geq \min\{\tau^+(M_1), \tau^+(M_2), \Lambda_n\}.$$

Note that these estimates on  $\tau^+$  would be trivial without Theorem 1.1.

**1.6. Surgery and bordism.** Performing surgery on a spin manifold is a special case of joining manifolds, this is discussed in more detail in Section 2.2. From Corollary 1.3 we get an inequality relating the  $\tau$ -invariant before and after surgery. For a compact spin manifold  $M$  of dimension  $n$  we define

$$\overline{\tau}^+(M) := \min\{\tau^+(M), \Lambda_n\}.$$

We also define

$$\bar{\tau}(M) := \min\{\tau^+(M), \tau^-(M), \Lambda_n\}.$$

If  $n \not\equiv 3 \pmod{4}$  then  $\bar{\tau}(M) = \bar{\tau}^+(M)$ . As noted before, all results for  $\bar{\tau}^+(M)$  also hold for  $\bar{\tau}^-(M) := \min\{\tau^-(M), \Lambda_n\}$ .

**Corollary 1.4.** *Assume that  $M$  is a spin manifold of dimension  $n$  and that  $N$  is obtained from  $M$  by a surgery of codimension  $n - k \geq 2$ . Then*

$$\tau^+(N) \geq \min\{\tau^+(M), \Lambda_{n,k}\} \geq \min\{\tau^+(M), \Lambda_n\}.$$

Corollary 1.4 tells us that

$$\tau^+(N) \geq \bar{\tau}^+(M), \quad \bar{\tau}(N) \geq \bar{\tau}(M).$$

Two compact spin manifolds  $M$  and  $N$  are spin bordant if there is a spin diffeomorphism from their disjoint union to the boundary of a spin manifold of one dimension higher, and this diffeomorphism respects the orientation of  $N$  and reverses that of  $M$ . This happens if and only if  $N$  can be obtained from  $M$  by a sequence of surgeries. To apply Corollary 1.4 we need to know when this sequence of surgeries can be chosen to include only surgeries of codimension at least two. The theory of handle decompositions of bordisms tells us that this can be done when  $N$  is connected, see [16, VII Theorem 3] for dimension 3, and [18, VIII Proposition 3.1] for higher dimensions.

**Corollary 1.5.** *Let  $M$  and  $N$  be spin bordant manifolds of dimension at least 3 and assume that  $N$  is connected. Then  $\bar{\tau}(N) \geq \bar{\tau}(M)$ . In particular, if  $M$  is also connected we have  $\bar{\tau}(N) = \bar{\tau}(M)$ .*

Corollary 1.5 can also be shown in dimension 2 with similar arguments [7, Theorem 1.3].

The spin bordism group  $\Omega_n^{\text{spin}}$  is the set of equivalence classes of spin bordant manifolds of dimension  $n$  with disjoint union as addition. Since every element in  $\Omega_n^{\text{spin}}$  can be represented by a connected manifold we obtain a well-defined map  $\bar{\tau} : \Omega_n^{\text{spin}} \rightarrow [0, \Lambda_n]$  which sends the equivalence class  $[M]$  of a connected spin manifold  $M$  to  $\bar{\tau}(M)$ .

**Corollary 1.6.** *There is a positive constant  $\varepsilon_n$  such that*

$$\tau^+(M) \in \{0\} \cup [\varepsilon_n, \lambda_{\min}^+(S^n, \sigma^n)].$$

for all spin manifolds  $M$  of dimension  $n$ .

*Proof.* The spin bordism group  $\Omega_n^{\text{spin}}$  is finitely generated [22, page 336]. This implies that the kernel of the map  $\alpha : \Omega_n^{\text{spin}} \rightarrow KO_n$  is also finitely generated. Let  $[N_1], \dots, [N_r]$  be generators of this kernel, we assume that the manifolds  $N_i$  are all connected. Since  $\tau(M) = 0$  if and only if  $\alpha(M) \neq 0$  we obtain the corollary for

$$\varepsilon_n := \min\{\Lambda_n, \bar{\tau}(N_1), \dots, \bar{\tau}(N_r)\}.$$

□

The  $\alpha$ -map is injective when  $n < 8$ , and then  $\varepsilon_n = \Lambda_n$ . We do not know whether there are  $n \in \mathbb{N}$  with  $\varepsilon_n < \Lambda_n$ . In other words, we do not know if there are  $n$ -dimensional manifolds  $M$  with  $0 < \tau^+(M) < \Lambda_n$ . If such manifolds exist, the following observations might be interesting.

First, if  $M$  is a spin manifold with  $\tau^+(M) < \Lambda_n$ , then it follows from Corollary 1.5 that the  $\sigma$ -constant of any manifold  $N$  spin bordant to  $M$  satisfies

$$\sigma(N) \leq \frac{4(n-1)}{n} \tau^+(M)^2.$$

For the next observation we define

$$S(t) := \{[M] \in \Omega_n^{\text{spin}} \mid \bar{\tau}(M) \geq t\}, \quad S^+(t) := \{[M] \in \Omega_n^{\text{spin}} \mid \bar{\tau}^+(M) \geq t\},$$

and

$$T(t) := \{[M] \in \Omega_n^{\text{spin}} \mid \bar{\tau}(M) > t\}, \quad T^+(t) := \{[M] \in \Omega_n^{\text{spin}} \mid \bar{\tau}^+(M) > t\}.$$

Obviously  $S(t) = S^+(t)$  and  $T(t) = T^+(t)$  in dimensions  $n \not\equiv 3 \pmod{4}$ .

**Corollary 1.7.**  *$S(t)$  is a subgroup of  $\Omega_n^{\text{spin}}$  for  $t \in [0, \Lambda_n]$  and  $T(t)$  is a subgroup of  $\Omega_n^{\text{spin}}$  for  $t \in [0, \Lambda_n)$ . If  $n \equiv 3 \pmod{4}$ , then  $S^+(t)$  and  $T^+(t)$  are submonoids.*

**Corollary 1.8.** *The values of  $\bar{\tau}$  cannot accumulate from above.*

*Proof.* Assume that  $t_i := \bar{\tau}(M_i)$ ,  $i \in \mathbb{N}$ , is a decreasing sequence of values of  $\bar{\tau}$  which converges to a limit  $t_\infty$ . We want to show that  $t_i = t_\infty$  for all but finitely many  $i$ .

We have  $S(t_i) \subset S(t_{i+1})$ , and hence  $\bigcup_i S(t_i) = T(t_\infty)$  is a subgroup of the finitely generated group  $\Omega_n^{\text{spin}}$ . It is thus finitely generated itself and we choose a finite set of generators. There must then be an  $I \in \mathbb{N}$  such that  $S(t_I)$  contains this finite set, and thus  $S(t_I) = T(t_\infty)$ . Hence  $[M_i] \in S(t_I)$  for all  $i$ , which implies  $t_i \geq t_I$ . We conclude that  $t_i = t_I = t_\infty$  for  $i \geq I$ .  $\square$

We do not know whether  $\bar{\tau}^+$  can accumulate from above in dimensions  $n \equiv 3 \pmod{4}$ .

**1.7. Variants of the results.** We already remarked earlier that if the alpha-genus  $\alpha(M)$  of a spin manifold  $M$  does not vanish, then the index theorem tells us that the kernel of  $D^g$  is non-trivial for any metric  $g$  on  $M$ , and hence  $\tau^+(M) = 0$ . For a connected spin manifold  $M$  the index theorem implies that the kernel of the Dirac operator has at least dimension

$$a(M) := \begin{cases} |\widehat{A}(M)|, & \text{if } n \equiv 0 \pmod{4}; \\ 1, & \text{if } n \equiv 1 \pmod{8} \text{ and } \alpha(M) \neq 0; \\ 2, & \text{if } n \equiv 2 \pmod{8} \text{ and } \alpha(M) \neq 0; \\ 0, & \text{otherwise.} \end{cases}$$

Let us modify the definition of  $\tau^+$  and use the  $k$ -th non-negative eigenvalue of the Dirac operator instead of the first one. The quantity thus obtained, denoted by  $\tau_k^+(M)$ , is zero if  $k \leq a(M)$ . It follows from [2] and [4] that  $\tau_{a(M)+1}^+(M) > 0$ . We expect that our methods generalize to this situation and yield similar surgery formulas for  $\tau_k^+$ .

## 2. PRELIMINARIES

**2.1. Notation for balls and neighbourhoods.** We write  $B^n(r)$  for the open standard ball of radius  $r$  around 0 in  $\mathbb{R}^n$ , and set  $B^n := B^n(1)$ . For a Riemannian manifold  $(M, g)$  we let  $B^g(p, r)$  denote the open ball of radius  $r$  around  $p \in M$ . If the Riemannian metric is clear from the context we will write  $B(p, r)$ . For a

Riemannian manifold  $(M, g)$  and a subset  $S \subset M$  we let  $U^g(S, r) := \bigcup_{x \in S} B^g(x, r)$  denote the  $r$ -neighbourhood of  $S$ . Again, if the Riemannian metric is clear from the context we abbreviate to  $U(S, r)$ .

**2.2. Joining manifolds along submanifolds.** We are now going to describe how two manifolds are joined along a common submanifold with trivialized normal bundle. Strictly speaking this is a differential topological construction, but since we work with Riemannian manifolds we will make the construction adapted to the Riemannian metrics and use distance neighbourhoods defined by the metrics etc.

Let  $(M_1, g_1)$  and  $(M_2, g_2)$  be complete Riemannian manifolds of dimension  $n$ . Let  $W$  be a compact manifold of dimension  $k$ , where  $0 \leq k \leq n$ . We assume that  $W$  is embedded in both  $M_1$  and  $M_2$  with trivializations of the normal bundle, we describe these embeddings as follows.

Let  $\bar{w}_i : W \times \mathbb{R}^{n-k} \rightarrow TM_i$ ,  $i = 1, 2$ , be smooth embeddings. We assume that  $\bar{w}_i$  restricted to  $W \times \{0\}$  maps to the zero section of  $TM_i$  (which we identify with  $M_i$ ) and thus gives an embedding  $W \rightarrow M_i$ . The image of this embedding is denoted by  $W'_i$ . Further we assume that  $\bar{w}_i$  restrict to linear isomorphisms  $\{p\} \times \mathbb{R}^{n-k} \rightarrow N_{\bar{w}_i(p,0)} W'_i$  for all  $p \in W$ , where  $NW'_i$  denotes the normal bundle of  $W'_i$  defined using  $g_i$ .

We set  $w_i := \exp^{g_i} \circ \bar{w}_i$ . For  $i = 1, 2$  this gives embeddings  $w_i : W \times B^{n-k}(R_{\max}) \rightarrow M_i$  for some  $R_{\max} > 0$ . We have  $W'_i = w_i(W \times \{0\})$  and we define the disjoint union

$$(M, g) := (M_1 \amalg M_2, g_1 \amalg g_2),$$

and

$$W' := W'_1 \amalg W'_2.$$

Let  $r_i$  be the function on  $M_i$  giving the distance to  $W'_i$ . Then  $r_1 \circ w_1(w, x) = r_2 \circ w_2(w, x) = |x|$  for  $w \in W$ ,  $x \in B^{n-k}(R_{\max})$ . Let  $r$  be the function on  $M$  defined by  $r(x) := r_i(x)$  for  $x \in M_i$ ,  $i = 1, 2$ . For  $0 < \varepsilon$  we set  $U_i(\varepsilon) := \{x \in M_i : r_i(x) < \varepsilon\}$  and  $U(\varepsilon) := U_1(\varepsilon) \cup U_2(\varepsilon)$ . For  $0 < \varepsilon < \theta$  we define

$$N_\varepsilon := (M_1 \setminus U_1(\varepsilon)) \cup (M_2 \setminus U_2(\varepsilon)) / \sim,$$

and

$$U_\varepsilon^N(\theta) := (U(\theta) \setminus U(\varepsilon)) / \sim$$

where  $\sim$  indicates that we identify  $x \in \partial U_1(\varepsilon)$  with  $w_2 \circ w_1^{-1}(x) \in \partial U_2(\varepsilon)$ . Hence

$$N_\varepsilon = (M \setminus U(\theta)) \cup U_\varepsilon^N(\theta).$$

We say that  $N_\varepsilon$  is obtained from  $M_1$ ,  $M_2$  (and  $\bar{w}_1$ ,  $\bar{w}_2$ ) by a connected sum along  $W$  with parameter  $\varepsilon$ .

The diffeomorphism type of  $N_\varepsilon$  is independent of  $\varepsilon$ , hence we will usually write  $N = N_\varepsilon$ . However, in some situations where dropping the index  $\varepsilon$  might cause ambiguities we will write  $N_\varepsilon$ . For example the function  $r : M_1 \amalg M_2 \rightarrow [0, \infty)$  also defines a continuous function  $r : N_\varepsilon \rightarrow [\varepsilon, \infty)$  whose definition depends on  $\varepsilon$ . We will also keep the  $\varepsilon$ -subscript for  $U_\varepsilon^N(\theta)$  as important estimates for spinors will be carried out on  $U_\varepsilon^N(\theta)$ . As the embeddings  $w_1$  and  $w_2$  preserve the spin structure, the manifold  $N$  carries a spin structure such that its restriction to  $(M_1 \setminus w_1(W \times B^{n-k})) \amalg (M_2 \setminus w_2(W \times B^{n-k}))$  coincides with the restriction of the given spin structure on  $M_1 \amalg M_2$ . If  $W$  is not connected, then this choice is not unique. The statements of our theorem hold for any such spin structure on  $N$ .

The surgery operation on a manifold is a special case of taking connected sum along a submanifold. Indeed, let  $M$  be a compact manifold of dimension  $n$  and let  $M_1 = M$ ,  $M_2 = S^n$ ,  $W = S^k$ . Let  $w_1 : S^k \times B^{n-k} \rightarrow M$  be an embedding defining a surgery and let  $w_2 : S^k \times B^{n-k} \rightarrow S^n$  be the standard embedding. Since  $S^n \setminus w_2(S^k \times B^{n-k})$  is diffeomorphic to  $B^{k+1} \times S^{n-k-1}$  we have in this situation that  $N$  is obtained from  $M$  using surgery on  $w_1$ , see [18, Section VI.9].

**2.3. Comparing spinors for different metrics.** Let  $M$  be a spin manifold of dimension  $n$  and let  $g, g'$  be Riemannian metrics on  $M$ . The goal of this paragraph is to identify the spinor bundles of  $(M, g)$  and  $(M, g')$  following Bourguignon and Gauduchon [9].

There exists a unique endomorphism  $b_{g'}^g$  of  $TM$  which is positive, symmetric with respect to  $g$ , and satisfies  $g(X, Y) = g'(b_{g'}^g X, b_{g'}^g Y)$  for all  $X, Y \in TM$ . This endomorphism maps  $g$ -orthonormal frames at a point to  $g'$ -orthonormal frames at the same point and we get a map  $b_{g'}^g : \text{SO}(M, g) \rightarrow \text{SO}(M, g')$  of  $\text{SO}(n)$ -principal bundles. If we assume that  $\text{Spin}(M, g)$  and  $\text{Spin}(M, g')$  are equivalent spin structures on  $M$  then the map  $b_{g'}^g$  lifts to a map  $\beta_{g'}^g$  of  $\text{Spin}(n)$ -principal bundles,

$$\begin{array}{ccc} \text{Spin}(M, g) & \xrightarrow{\beta_{g'}^g} & \text{Spin}(M, g') \\ \downarrow & & \downarrow \\ \text{SO}(M, g) & \xrightarrow{b_{g'}^g} & \text{SO}(M, g') \end{array} .$$

From this we get a map between the spinor bundles  $\Sigma^g M$  and  $\Sigma^{g'} M$  denoted by the same symbol and defined by

$$\begin{aligned} \beta_{g'}^g : \Sigma^g M = \text{Spin}(M, g) \times_{\rho} \Sigma_n &\rightarrow \text{Spin}(M, g') \times_{\rho} \Sigma_n = \Sigma^{g'} M, \\ \psi = [s, \varphi] &\mapsto [\beta_{g'}^g s, \varphi] = \beta_{g'}^g \psi, \end{aligned}$$

where  $(\rho, \Sigma_n)$  is the complex spinor representation, and where  $[s, \varphi] \in \text{Spin}(M, g) \times_{\rho} \Sigma_n$  denotes the equivalence class of  $(s, \varphi) \in \text{Spin}(M, g) \times \Sigma_n$  for the equivalence relation given by the action of  $\text{Spin}(n)$ . The map  $\beta_{g'}^g$  of Hermitian vector bundles is fiberwise an isometry.

We define the Dirac operator  ${}^g D^{g'}$  acting on sections of the spinor bundle for  $g$  by

$${}^g D^{g'} := (\beta_{g'}^g)^{-1} \circ D^{g'} \circ \beta_{g'}^g.$$

In [9, Theorem 20] the operator  ${}^g D^{g'}$  is computed in terms of  $D^g$  and some extra terms which are small if  $g$  and  $g'$  are close. Formulated in a way convenient for us the relationship is

$${}^g D^{g'} \psi = D^g \psi + A_{g'}^g(\nabla^g \psi) + B_{g'}^g(\psi), \quad (3)$$

where  $A_{g'}^g \in \text{hom}(T^*M \otimes \Sigma^g M, \Sigma^g M)$  satisfies

$$|A_{g'}^g| \leq C|g - g'|_g, \quad (4)$$

and  $B_{g'}^g \in \text{hom}(\Sigma^g M, \Sigma^g M)$  satisfies

$$|B_{g'}^g| \leq C(|g - g'|_g + |\nabla^g(g - g')|_g) \quad (5)$$

for some constant  $C$ .



In the special case that  $g'$  and  $g$  are conformal with  $g' = F^2g$  for a positive smooth function  $F$  the formula simplifies considerably, and one obtains

$${}^gD^{g'}(F^{-\frac{n-1}{2}}\psi) = F^{-\frac{n+1}{2}}D^g\psi, \quad (6)$$

see for instance [15, 8].

**2.4. Regularity results.** By standard elliptic theory we have the following lemma (see for example [3, Chapter 3] where the corresponding results of [11] are adapted to the Dirac operator).

**Lemma 2.1.** *Let  $(V, g)$  be a Riemannian spin manifold and  $\Omega \subset V$  an open set with compact closure in  $V$ . Let also  $r \in (1, \infty)$ . Then there is a constant  $C$  so that*

$$\int_{\Omega} |\nabla^g \varphi|^r dv^g \leq C \left( \int_{\Omega} |D^g \varphi|^r dv^g + \int_{\Omega} |\varphi|^r dv^g \right) \quad (7)$$

for all  $\varphi \in \Gamma(\Sigma^g \Omega)$  which are of class  $C^1$  and compactly supported in  $\Omega$ .

For a compact Riemannian manifold with invertible Dirac operator we have the following special case.

**Lemma 2.2.** *Let  $(V, g)$  be a compact Riemannian spin manifold such that  $D^g$  is invertible. Then there exists a constant  $C$  such that*

$$\int_V |\nabla^g \varphi|^{\frac{2n}{n+1}} dv^g \leq C \int_V |D^g \varphi|^{\frac{2n}{n+1}} dv^g \quad (8)$$

for all  $\varphi \in \Gamma(\Sigma^g V)$  of class  $C^1$ .

**2.5. The associated variational problem.** Let  $(M, g)$  be a compact spin manifold of dimension  $n$  with  $\ker D^g = \{0\}$ . We define the functional  $J^g$  acting on smooth spinor fields  $\psi \in \Gamma(\Sigma^g M)$  by

$$J^g(\psi) := \frac{\left( \int_M |D\psi|^{\frac{2n}{n+1}} dv^g \right)^{\frac{n+1}{n}}}{\int_M \langle D\psi, \psi \rangle dv^g},$$

whenever the denominator is non-zero. Using techniques from [20] it was proved in [2] that

$$\lambda_{\min}^+(M, g) = \inf_{\psi} J^g(\psi), \quad (9)$$

where the infimum is taken over the set of smooth spinor fields satisfying

$$\int_M \langle D\psi, \psi \rangle dv^g > 0.$$

If  $g$  and  $\tilde{g} = F^2g$  are conformal metrics on  $M$  and if  $J^g$  and  $J^{\tilde{g}}$  are the associated functionals, then by Relation (6) one computes that

$$J^{\tilde{g}}(F^{-\frac{n-1}{2}}\psi) = J^g(\psi) \quad (10)$$

for smooth  $\psi \in \Gamma(\Sigma^g M)$ .

The following result gives a universal upper bound on  $\lambda_{\min}^+(M, g)$ .

**Proposition 2.3.** *Let  $(M, g)$  be a compact spin manifold of dimension  $n \geq 2$ . Then*

$$\lambda_{\min}^+(M, g) \leq \lambda_{\min}^+(S^n, \sigma^n) = \frac{n}{2} \omega_n^{1/n}, \quad (11)$$

where  $\omega_n$  is the volume of  $(S^n, \sigma^n)$ .

Proposition 2.3 was proven for  $n \geq 3$  in [2] using geometric methods. In the case  $n = 2$  the article [2] only provides a proof if  $\ker D = \{0\}$ . Another method that yields the proposition in full generality is to construct for any  $p \in M$  and  $\varepsilon > 0$  a suitable test spinor field  $\psi_\varepsilon$  supported in  $B^g(p, \varepsilon)$  satisfying  $J^g(\psi_\varepsilon) \leq \lambda_{\min}^+(S^n, \sigma^n) + o(\varepsilon)$ , see [6] for details.

If Inequality (11) holds strictly then one can show that the infimum in Equation (9) is attained by a spinor field  $\varphi$ . The following theorem will be a central ingredient in the proof of Theorem 1.2.

**Theorem 2.4** ([1, 3]). *Let  $(M, g)$  be a compact spin manifold of dimension  $n$  for which Inequality (11) holds strictly. Then there exists a spinor field  $\varphi \in C^{2,\alpha}(\Sigma M) \cap C^\infty(\Sigma M \setminus \varphi^{-1}(0))$  where  $\alpha \in (0, 1) \cap (0, 2/(n-1)]$  such that  $\|\varphi\|_{L^{\frac{2n}{n-1}}(M)} = 1$  and*

$$D\varphi = \lambda_{\min}^+(M, g)|\varphi|^{\frac{2}{n-1}}\varphi.$$

Furthermore the infimum in the definition of  $\lambda_{\min}^+(M, g)$  is attained by the generalized conformal metric  $\tilde{g} = |\varphi|^{4/(n-1)}g$ , see [1] for details.

### 3. PREPARATIONS FOR PROOFS

**3.1. Removal of singularities.** The following theorem gives a condition for when singularities of solutions to Dirac equations can be removed.

**Theorem 3.1.** *Let  $(V, g)$  be a (not necessarily complete) Riemannian spin manifold and let  $S$  be a compact submanifold of  $V$  of codimension  $m \geq 2$ . Assume that  $\varphi \in L^p(\Sigma(V \setminus S))$ ,  $p \geq m/(m-1)$ , satisfies the equation*

$$D\varphi = \rho$$

*weakly on  $V \setminus S$  where  $\rho \in L^1(\Sigma(V \setminus S)) = L^1(\Sigma V)$ . Then this equation holds weakly on  $V$ . In particular the singular support of the distribution  $D\varphi$  is empty.*

*Proof.* Let  $\psi$  be a smooth compactly supported spinor. We have to show that

$$\int_V \langle \varphi, D\psi \rangle dv = \int_V \langle \rho, \psi \rangle dv. \quad (12)$$

Recall that for  $\varepsilon > 0$  we denote the set of points in  $V$  of distance less than  $\varepsilon$  to  $S$  by  $U(S, \varepsilon)$ . We choose a smooth cut-off function  $\chi_\varepsilon : V \rightarrow [0, 1]$  with support in  $U(S, 2\varepsilon)$ ,  $\chi_\varepsilon = 1$  on  $U(S, \varepsilon)$ , and  $|\text{grad}\chi_\varepsilon| \leq 2/\varepsilon$ . We then have

$$\begin{aligned} \int_V \langle \varphi, D\psi \rangle dv - \int_V \langle \rho, \psi \rangle dv &= \int_V \langle \varphi, D((1 - \chi_\varepsilon)\psi + \chi_\varepsilon\psi) \rangle dv - \int_V \langle \rho, \psi \rangle dv \\ &= \int_V \langle D\varphi, (1 - \chi_\varepsilon)\psi \rangle dv + \int_V \langle \varphi, \chi_\varepsilon D\psi \rangle dv \\ &\quad + \int_V \langle \varphi, \text{grad}\chi_\varepsilon \cdot \psi \rangle dv - \int_V \langle \rho, \psi \rangle dv \\ &= - \int_V \langle \rho, \chi_\varepsilon\psi \rangle dv + \int_V \langle \varphi, \chi_\varepsilon D\psi \rangle dv \\ &\quad + \int_V \langle \varphi, \text{grad}\chi_\varepsilon \cdot \psi \rangle dv, \end{aligned}$$

where  $D\varphi = \rho$  is used in the last equality. Let  $q$  be related to  $p$  via  $1/q + 1/p = 1$ . It follows that

$$\begin{aligned} \left| \int_V \langle \varphi, D\psi \rangle dv - \int_V \langle \rho, \psi \rangle dv \right| &\leq \left( \sup_{U(S, 2\varepsilon)} |\psi| \right) \int_{U(S, 2\varepsilon)} |\rho| dv \\ &\quad + \left( \sup_{U(S, 2\varepsilon)} |D\psi| + \frac{2}{\varepsilon} \sup_{U(S, 2\varepsilon)} |\psi| \right) \int_{U(S, 2\varepsilon)} |\varphi| dv \\ &\leq o(1) + \frac{C}{\varepsilon} \|\varphi\|_{L^p(U(S, 2\varepsilon))} \text{Vol}(U(S, 2\varepsilon))^{1/q} \\ &\leq o(1) + C \|\varphi\|_{L^p(U(S, 2\varepsilon))} \varepsilon^{(m/q)-1}, \end{aligned}$$

where  $o(1)$  denotes a term tending to 0 as  $\varepsilon \rightarrow 0$ . Since  $p \geq m/(m-1)$  is equivalent to  $m/q \geq 1$  we see that (12) holds.  $\square$

Applying Theorem 3.1 to the non-linear Dirac equation in Theorem 2.4 we get the following corollary.

**Corollary 3.2.** *Let  $V$  and  $S$  be as in Theorem 3.1. Then any  $L^p$ -solution,  $p = 2n/(n-1)$ , of*

$$D\varphi = \lambda |\varphi|^{p-2} \varphi \tag{13}$$

on  $V \setminus S$  is also a weak  $L^p$ -solution of (13) on  $V$ .

**3.2. Limit spaces and limit solutions.** In the proofs of the main theorems we will construct limit solutions of a Dirac equation on certain limit spaces. For this we need the following two lemmas. In the statement of these results, in order to simplify the notations, we write  $\alpha \rightarrow 0$  instead of  $\alpha_i \rightarrow 0$  as  $i \rightarrow \infty$  when  $(\alpha_i)_{i \in \mathbb{N}}$  is a sequence of positive numbers converging to 0. In the same way, the subsequences of  $(\alpha_i)$  will also be denoted by  $(\alpha)$ .

**Lemma 3.3.** *Let  $V$  be an  $n$ -dimensional manifold. Let  $(p_\alpha)$  be a sequence of points in  $V$  which converges to a point  $p$  as  $\alpha \rightarrow 0$ . Let  $(\gamma_\alpha)$  be a sequence of metrics defined on a neighbourhood  $O$  of  $p$  which converges to a metric  $\gamma_0$  in the  $C^2(O)$ -topology. Finally, let  $(b_\alpha)$  be a sequence of positive real numbers such that  $\lim_{\alpha \rightarrow 0} b_\alpha = \infty$ . Then for  $r > 0$  there exists for  $\alpha$  small enough a diffeomorphism*

$$\Theta_\alpha : B^n(r) \rightarrow B^{\gamma_\alpha}(p_\alpha, b_\alpha^{-1}r)$$

with  $\Theta_\alpha(0) = p_\alpha$  such that the metric  $\Theta_\alpha^*(b_\alpha^2 \gamma_\alpha)$  tends to the Euclidean metric  $\xi^n$  in  $C^1(B^n(r))$ .

*Proof.* Denote by  $\exp_{p_\alpha}^{\gamma_\alpha} : U_\alpha \rightarrow O_\alpha$  the exponential map at the point  $p_\alpha$  defined with respect to the metric  $\gamma_\alpha$ . Here  $O_\alpha$  is a neighbourhood of  $p_\alpha$  in  $V$  and  $U_\alpha$  is a neighbourhood of the origin in  $\mathbb{R}^n$ . We set

$$\Theta_\alpha : B^n(r) \ni x \mapsto \exp_{p_\alpha}^{\gamma_\alpha}(b_\alpha^{-1}x) \in B^{\gamma_\alpha}(p_\alpha, b_\alpha^{-1}r).$$

It is easily checked that  $\Theta_\alpha$  is the desired diffeomorphism.  $\square$

**Lemma 3.4.** *Let  $V$  an  $n$ -dimensional spin manifold. Let  $(g_\alpha)$  be a sequence of metrics which converges to a metric  $g$  in  $C^1$  on all compact sets  $K \subset V$  as  $\alpha \rightarrow 0$ . Assume that  $(U_\alpha)$  is an increasing sequence of subdomains of  $V$  such that  $\cup_\alpha U_\alpha =$*

$V$ . Let  $\psi_\alpha \in \Gamma(\Sigma^{g_\alpha} U_\alpha)$  be a sequence of spinors of class  $C^1$  such that  $\|\psi_\alpha\|_{L^\infty(U_\alpha)} \leq C$  where  $C$  does not depend on  $\alpha$ , and

$$D^{g_\alpha} \psi_\alpha = \lambda_\alpha |\psi_\alpha|^{\frac{2}{n-1}} \psi_\alpha \quad (14)$$

where the  $\lambda_\alpha$  are positive numbers which tend to  $\bar{\lambda} \geq 0$ . Then there exists a spinor  $\psi \in \Gamma(\Sigma^g V)$  of class  $C^1$  such that

$$D^g \psi = \bar{\lambda} |\psi|^{\frac{2}{n-1}} \psi \quad (15)$$

on  $V$  and a subsequence of  $(\beta_{g_\alpha}^{g_\alpha} \psi_\alpha)$  tends to  $\psi$  in  $C^0(K)$  for any compact set  $K \subset V$ . In particular

$$\|\psi\|_{L^\infty(K)} = \lim_{\alpha \rightarrow 0} \|\psi_\alpha\|_{L^\infty(K)}, \quad (16)$$

and

$$\int_K |\psi|^r dv^g = \lim_{\alpha \rightarrow 0} \int_K |\psi_\alpha|^r dv^{g_\alpha} \quad (17)$$

for any compact set  $K$  and any  $r \geq 1$ .

*Proof.* Let  $K$  be a compact subset of  $V$  and let  $\Omega$  be an open set in  $V$  with compact closure such that  $K \subset \Omega$ . Let  $\chi \in C^\infty(V)$  with  $0 \leq \chi \leq 1$  be compactly supported in  $\Omega$  and satisfy  $\chi = 1$  on a neighbourhood  $\tilde{\Omega}$  of  $K$ . Set  $\varphi_\alpha = (\beta_{g_\alpha}^{g_\alpha})^{-1} \psi_\alpha$ . Using Equations (14) and (3) we get

$$D^g(\chi \varphi_\alpha) = \text{grad}^g \chi \cdot \varphi_\alpha + \chi \lambda_\alpha |\varphi_\alpha|^{\frac{2}{n-1}} \varphi_\alpha - \chi A_{g_\alpha}^g(\nabla^g \varphi_\alpha) - \chi B_{g_\alpha}^g(\varphi_\alpha). \quad (18)$$

Using the fact that  $|a + b + c|^r \leq 3^r(|a|^r + |b|^r + |c|^r)$  for  $a, b, c \in \mathbb{R}$ ,  $r \geq 1$ , we see that

$$\begin{aligned} |D^g(\chi \varphi_\alpha)|^r &\leq 3^r \left( |\text{grad}^g \chi \cdot \varphi_\alpha + \chi \lambda_\alpha |\varphi_\alpha|^{\frac{2}{n-1}} \varphi_\alpha|^r \right. \\ &\quad \left. + |\chi A_{g_\alpha}^g(\nabla^g \varphi_\alpha)|^r + |\chi B_{g_\alpha}^g(\varphi_\alpha)|^r \right) \end{aligned}$$

for  $r \geq 1$ . Since  $\|\varphi_\alpha\|_{L^\infty(V)} = \|\psi_\alpha\|_{L^\infty(V)} \leq C$  we have

$$|\text{grad}^g \chi \cdot \varphi_\alpha + \chi \lambda_\alpha |\varphi_\alpha|^{\frac{2}{n-1}} \varphi_\alpha|^r \leq C.$$

By Relations (4) and (5), and since  $\lim_{\alpha \rightarrow 0} \|g_\alpha - g\|_{C^1(\Omega)} = 0$ , we get

$$\begin{aligned} |\chi A_{g_\alpha}^g(\nabla^g \varphi_\alpha)|^r + |\chi B_{g_\alpha}^g(\varphi_\alpha)|^r &\leq o(1) (|\nabla^g(\chi \varphi_\alpha)|^r + |\text{grad}^g \chi \cdot \varphi_\alpha|^r + |\chi \varphi_\alpha|^r) \\ &\leq o(1) (|\nabla^g(\chi \varphi_\alpha)|^r + C), \end{aligned}$$

where  $o(1)$  tends to 0 with  $\alpha$ . It follows that

$$|D^g(\chi \varphi_\alpha)|^r \leq C + o(1) |\nabla^g(\chi \varphi_\alpha)|^r.$$

Setting  $\varphi = \chi \varphi_\alpha$  in Inequality (7) and again using that  $\|\varphi_\alpha\|_{L^\infty(\Omega)}$  is uniformly bounded we get that

$$\int_\Omega |\nabla^g(\chi \varphi_\alpha)|^r dv^g \leq C + o(1) \int_\Omega |\nabla^g(\chi \varphi_\alpha)|^r dv^g.$$

In particular  $(\chi \varphi_\alpha)$  is bounded in  $H_0^{1,r}(\Omega)$ . Let  $a \in (0, 1)$ . By the Sobolev Embedding Theorem this implies that a subsequence of  $(\chi \varphi_\alpha)$  converges in  $C^{0,a}(\Omega)$  to  $\psi_K \in \Gamma(\Sigma^{g_\alpha} \Omega)$  of class  $C^{0,a}$ . We take the inner product of (18) with a smooth

spinor  $\tilde{\varphi}$  which is compactly supported in  $\tilde{\Omega}$  and integrate over  $\Omega$ . Since  $\chi = 1$  on the support of  $\tilde{\varphi}$  the result is

$$\begin{aligned} \int_{\Omega} \langle \varphi_{\alpha}, D^g \tilde{\varphi} \rangle dv^g &= \int_{\Omega} \lambda_{\alpha} |\varphi_{\alpha}|^{\frac{2}{n-1}} \langle \varphi_{\alpha}, \tilde{\varphi} \rangle dv^g \\ &\quad - \int_{\Omega} \langle A_{g_{\alpha}}^g(\nabla^g \varphi_{\alpha}), \tilde{\varphi} \rangle dv^g - \int_{\Omega} \langle B_{g_{\alpha}}^g(\varphi_{\alpha}), \tilde{\varphi} \rangle dv^g. \end{aligned}$$

Taking the limit  $\alpha \rightarrow 0$  and again using (4) and (5) we get

$$\int_{\Omega} \langle \psi_K, D^g \tilde{\varphi} \rangle dv^g = \int_{\Omega} \bar{\lambda} |\psi_K|^{\frac{2}{n-1}} \langle \psi_K, \tilde{\varphi} \rangle dv^g.$$

Hence,  $\psi_K$  satisfies Equation (15) weakly on  $K$ . By standard regularity theorems we conclude that  $\psi_K \in C^1(K)$ .

Now we choose an increasing sequence of compact sets  $K_m$  such that  $\cup_m K_m = V$ . Using the above arguments and taking successive subsequences it follows that  $(\varphi_{\alpha})$  converge to spinor fields  $\psi_m$  on  $K_m$  with  $\psi_m|_{K_{m-1}} = \psi_{m-1}$ . We define  $\psi$  on  $V$  by  $\psi := \psi_m$  on  $K_m$ . By taking a diagonal subsequence of we get that  $(\varphi_{\alpha})$  tends to  $\psi$  in  $C^0$  on any compact set  $K \subset V$ .

The relations (16) and (17) follow immediately since  $\beta_{g_{\alpha}}^g$  is an isometry, since  $\varphi_{\alpha} = (\beta_{g_{\alpha}}^g)^{-1} \psi_{\alpha}$ , and since  $(g_{\alpha})$  (resp.  $(\varphi_{\alpha})$ ) tends to  $g$  (resp.  $\psi$ ) in  $C^0$  on  $K$ . This ends the proof of Lemma 3.4.  $\square$

**3.3. Dirac spectral bounds on products with spheres.** In the following lemma we assume (in the case  $m = 1$ ) that  $S^1$  carries the spin structure which is obtained by restricting the unique spin structure on the  $B^2$  to the boundary. The proof is a simple application of the formula for the squared Dirac operator on a product manifold together with the lower bound of its spectrum on the standard sphere.

**Lemma 3.5.** *Let  $(V, g)$  be a complete Riemannian spin manifold. Then any  $L^2$ -spinor  $\psi$  on  $(V \times S^m, g + \sigma^m)$  satisfies*

$$\int_{V \times S^m} |D\psi|^2 dv^{g+\sigma^m} \geq \frac{m^2}{4} \int_{V \times S^m} |\psi|^2 dv^{g+\sigma^m}.$$

**3.4. Approximation by local product metrics.** In this paragraph we will see how to change the metrics  $g_i$  to product form  $g_i = h_i + dr_i^2 + r_i^2 \sigma^{n-k-1}$  in a neighbourhood of  $W'_i$  in  $M_i$  without changing  $\lambda_{\min}^+(M_i, g_i)$  much.

**Lemma 3.6.** *Let  $(V, g)$  be a compact Riemannian manifold of dimension  $n$  and let  $S$  be a closed submanifold of dimension  $k$ , where  $0 \leq k \leq n - 2$ . Assume that a trivialization of the normal bundle of  $S$  is given and assume that  $D^g$  is invertible. Then there exists a sequence  $(\varepsilon_i)_{i \in \mathbb{N}}$  of positive real numbers converging to 0 and a sequence  $(g_{\varepsilon_i})$  of metrics on  $V$  such that*

$$\lim_{i \rightarrow \infty} \lambda_{\min}^+(V, g_{\varepsilon_i}) = \lambda_{\min}^+(V, g)$$

and

$$g_{\varepsilon_i} = h + dr^2 + r^2 \sigma^{n-k-1}$$

on  $U^g(S, \varepsilon_i)$ . Here  $h$  is the restriction of the metric  $g$  to  $S$  and  $r(x) = d^g(S, x)$ .

*Proof.* Using the trivialization of the normal bundle we identify a neighbourhood of  $S$  with  $S \times B^{n-k}(R_{\max})$  as described in Section 2.2. In this neighbourhood we define the metric  $\bar{g} := h + dr^2 + r^2 \sigma^{n-k-1}$ . Recall that  $U^g(S, \varepsilon)$  denotes the set

of points  $x \in V$  such that  $r(x) < \varepsilon$  and let  $\chi_\varepsilon \in C^\infty(M)$ ,  $0 \leq \chi \leq 1$ , be a cut-off function such that  $\chi = 1$  on  $U^g(S, \varepsilon)$ ,  $\chi = 0$  on  $M \setminus U^g(S, 2\varepsilon)$ , and  $|d\chi_\varepsilon| \leq 2/\varepsilon$ . We define

$$g_\varepsilon := \chi_\varepsilon \bar{g} + (1 - \chi_\varepsilon)g.$$

Then  $g_\varepsilon$  has product form on  $U^g(S, \varepsilon)$ . For convenience we introduce the notation  $\lambda_\varepsilon := \lambda_{\min}^+(V, g_\varepsilon)$  and  $\lambda := \lambda_{\min}^+(V, g)$ . Let  $(\varepsilon_i)_{i \in \mathbb{N}}$  be a sequence of positive numbers tending to 0 such that the limit  $\lim_{i \rightarrow \infty} \lambda_{\varepsilon_i}$  exists. In the following, we write  $\varepsilon \rightarrow 0$  instead of  $\varepsilon_i \rightarrow 0$  as  $i \rightarrow \infty$ . In the same way,  $(\varepsilon)$  will denote the successive subsequences of  $(\varepsilon_i)$  we will need. With this notations, let  $\bar{\lambda} := \lim_{\varepsilon \rightarrow 0} \lambda_\varepsilon$  which exists after possibly taking a subsequence.

We begin by proving that

$$\bar{\lambda} \leq \lambda, \quad (19)$$

which is the simpler part of the proof. Let  $J := J^g$  and  $J_\varepsilon := J^{g_\varepsilon}$  be the functionals associated to  $g$  and  $g_\varepsilon$ , and let  $\delta > 0$  be a small number. We set  $\chi'_\varepsilon := 1 - \chi_{2\varepsilon}$ , so that  $\chi'_\varepsilon = 1$  on  $V \setminus U^g(S, 4\varepsilon)$ ,  $\chi'_\varepsilon = 0$  on  $U^g(S, 2\varepsilon)$ , and  $|d\chi'_\varepsilon| \leq 1/\varepsilon$ . We see that  $g = g_\varepsilon$  on the support of  $\eta'_\varepsilon$ . Let  $\psi$  be a smooth spinor such that  $J(\psi) \leq \lambda + \delta$ . We then have

$$\int_V \langle D^g(\chi'_\varepsilon \psi), \chi'_\varepsilon \psi \rangle dv^g = \int_V \chi_\varepsilon'^2 \langle D^g \psi, \psi \rangle dv^g + \int_V \langle \text{grad}^g \chi'_\varepsilon \cdot \psi, \chi'_\varepsilon \psi \rangle dv^g.$$

Since the last term here is purely imaginary we obtain

$$\lim_{\varepsilon \rightarrow 0} \int_V \langle D^g(\chi'_\varepsilon \psi), \chi'_\varepsilon \psi \rangle dv^g = \lim_{\varepsilon \rightarrow 0} \text{Re} \int_V \chi_\varepsilon'^2 \langle D^g \psi, \psi \rangle dv^g = \int_V \langle D^g \psi, \psi \rangle dv^g. \quad (20)$$

We compute

$$\begin{aligned} \int_V |D^g(\chi'_\varepsilon \psi)|^{\frac{2n}{n+1}} dv^g &= \int_{V \setminus U^g(S, 4\varepsilon)} |D^g \psi|^{\frac{2n}{n+1}} dv^g \\ &\quad + \int_{U^g(S, 4\varepsilon) \setminus U^g(S, 2\varepsilon)} |\text{grad}^g \chi'_\varepsilon \cdot \psi + \chi'_\varepsilon D^g \psi|^{\frac{2n}{n+1}} dv^g. \end{aligned} \quad (21)$$

Using the fact that  $|a + b|^{\frac{2n}{n+1}} \leq 2^{\frac{2n}{n+1}} (|a|^{\frac{2n}{n+1}} + |b|^{\frac{2n}{n+1}})$  for  $a, b \in \mathbb{R}$  we have

$$\begin{aligned} |\text{grad}^g \chi'_\varepsilon \cdot \psi + \chi'_\varepsilon D^g \psi|^{\frac{2n}{n+1}} &\leq 2^{\frac{2n}{n+1}} \left( |\text{grad}^g \chi'_\varepsilon|^{\frac{2n}{n+1}} |\psi|^{\frac{2n}{n+1}} + |\chi'_\varepsilon|^{\frac{2n}{n+1}} |D^g \psi|^{\frac{2n}{n+1}} \right) \\ &\leq 2^{\frac{2n}{n+1}} \left( C_1 \varepsilon^{-\frac{2n}{n+1}} + C_2 \right), \end{aligned}$$

where  $C_1$  and  $C_2$  are bounds on  $|\psi|$  and  $|D\psi|$ . Since  $\text{Vol}(U^g(S, 4\varepsilon) \setminus U^g(S, 2\varepsilon)) \leq C\varepsilon^{n-k} \leq C\varepsilon^2$  it follows that

$$\lim_{\varepsilon \rightarrow 0} \int_{U^g(S, 4\varepsilon) \setminus U^g(S, 2\varepsilon)} |\text{grad}^g \chi'_\varepsilon \cdot \psi + \chi'_\varepsilon D^g \psi|^{\frac{2n}{n+1}} dv^g = 0.$$

It is clear that  $\lim_{\varepsilon \rightarrow 0} \int_{V \setminus U^g(S, 4\varepsilon)} |D^g \psi|^{\frac{2n}{n+1}} dv^g = \int_V |D^g \psi|^{\frac{2n}{n+1}} dv^g$  so Equation (21) tells us that

$$\lim_{\varepsilon \rightarrow 0} \int_V |D^g(\chi'_\varepsilon \psi)|^{\frac{2n}{n+1}} dv^g = \int_V |D^g \psi|^{\frac{2n}{n+1}} dv^g.$$

Together with Equation (20) this proves that  $\lim_{\varepsilon \rightarrow 0} J(\chi'_\varepsilon \psi) = J(\psi) \leq \lambda + \delta$ . Since  $g_\varepsilon = g$  on the support of  $\chi'_\varepsilon \psi$ , we have  $J_\varepsilon(\chi'_\varepsilon \psi) = J(\chi'_\varepsilon \psi)$ . Relation (19) now follows since  $\lambda_\varepsilon \leq J_\varepsilon(\chi'_\varepsilon \psi)$  and  $\delta$  is arbitrary.

The second and harder part of the proof is to show that

$$\bar{\lambda} \geq \lambda. \quad (22)$$

From Proposition 2.3 we know that  $\lambda_\varepsilon \leq \lambda_{\min}^+(S^n, \sigma^n)$ ,  $\bar{\lambda} \leq \lambda_{\min}^+(S^n, \sigma^n)$ , and  $\lambda \leq \lambda_{\min}^+(S^n, \sigma^n)$ . Inequality (22) is obvious if  $\lambda = \lambda_{\min}^+(S^n, \sigma^n)$ . Hence we will assume  $\lambda_\varepsilon < \lambda_{\min}^+(S^n, \sigma^n)$  for a sequence  $\varepsilon \rightarrow 0$ . As the Dirac operator is invertible we know that (8) holds. By Theorem 2.4 there exists for all  $\varepsilon$  spinor fields  $\psi_\varepsilon \in \Gamma(\Sigma^{g_\varepsilon} v)$  of class  $C^1$  such that

$$D^{g_\varepsilon} \psi_\varepsilon = \lambda_\varepsilon |\psi_\varepsilon|^{\frac{2}{n-1}} \psi_\varepsilon, \quad (23)$$

and

$$\int_V |\psi_\varepsilon|^{\frac{2n}{n-1}} dv^{g_\varepsilon} = 1. \quad (24)$$

Define  $\varphi_\varepsilon = (\beta_{g_\varepsilon}^g)^{-1} \psi_\varepsilon$ . Since  $g_\varepsilon \rightarrow g$  it is easily seen that the sequence  $(\varphi_\varepsilon)$  is bounded in  $L^{\frac{2n}{n-1}}(V, g)$ . By (3) and (23) we have

$$D^g \varphi_\varepsilon = \lambda_\varepsilon |\varphi_\varepsilon|^{\frac{2}{n-1}} \varphi_\varepsilon - A_{g_\varepsilon}^g(\nabla^g \varphi_\varepsilon) - B_{g_\varepsilon}^g(\varphi_\varepsilon), \quad (25)$$

together with  $|a + b + c|^{\frac{2n}{n+1}} \leq 3^{\frac{2n}{n+1}} (|a|^{\frac{2n}{n+1}} + |b|^{\frac{2n}{n+1}} + |c|^{\frac{2n}{n+1}})$  for  $a, b, c \in \mathbb{R}$  this implies

$$|D^g \varphi_\varepsilon|^{\frac{2n}{n+1}} \leq C \left( \lambda_\varepsilon^{\frac{2n}{n+1}} |\varphi_\varepsilon|^{\frac{2n}{n-1}} + |A_{g_\varepsilon}^g(\nabla^g \varphi_\varepsilon)|^{\frac{2n}{n+1}} + |B_{g_\varepsilon}^g(\varphi_\varepsilon)|^{\frac{2n}{n+1}} \right). \quad (26)$$

We also have

$$|A_{g_\varepsilon}^g(\nabla^g \varphi_\varepsilon)| \leq \|g - g_\varepsilon\|_{C^0(V)} |\nabla^g \varphi_\varepsilon| \leq C\varepsilon |\nabla^g \varphi_\varepsilon|, \quad (27)$$

and

$$|B_{g_\varepsilon}^g(\varphi_\varepsilon)| \leq \|g - g_\varepsilon\|_{C^1(V)} |\varphi_\varepsilon| \leq C |\varphi_\varepsilon|. \quad (28)$$

Indeed, since  $g$  and  $g_\varepsilon$  coincide on  $S$ , there exists a constant  $C$  so that  $\|g - g_\varepsilon\|_{B^g(V, \varepsilon)} \leq C\varepsilon$ . Together with the fact that  $|d\chi_\varepsilon| \leq 2/\varepsilon$  and using the definition of  $g_\varepsilon$ , this immediately implies that  $\|g - g_\varepsilon\|_{C^1(V)} \leq C$ . Using Relation (8) and integrating (26) we find that

$$\begin{aligned} \int_V |\nabla^g \varphi_\varepsilon|^{\frac{2n}{n+1}} dv^g &\leq C \left( \lambda_\varepsilon^{\frac{2n}{n+1}} \int_V |\varphi_\varepsilon|^{\frac{2n}{n-1}} dv^g + \varepsilon^{\frac{2n}{n+1}} \int_V |\nabla^g \varphi_\varepsilon|^{\frac{2n}{n+1}} dv^g \right. \\ &\quad \left. + \int_V |B_{g_\varepsilon}^g(\varphi_\varepsilon)|^{\frac{2n}{n+1}} dv^g \right). \end{aligned}$$

As  $g$  and  $g_\varepsilon$  coincide on  $V \setminus B^g(S, 2\varepsilon)$  we conclude that  $B_{g_\varepsilon}^g(\varphi_\varepsilon) = 0$  on this set. Together with (28) we have

$$\begin{aligned} \int_V |B_{g_\varepsilon}^g(\varphi_\varepsilon)|^{\frac{2n}{n+1}} dv^g &\leq C \int_{B^g(S, 2\varepsilon)} |\varphi_\varepsilon|^{\frac{2n}{n+1}} dv^g \\ &\leq C \text{Vol}(B^g(S, 2\varepsilon))^{\frac{2}{n+1}} \left( \int_{B^g(S, 2\varepsilon)} |\varphi_\varepsilon|^{\frac{2n}{n-1}} dv^g \right)^{\frac{n-1}{n+1}} \\ &= o(1), \end{aligned}$$

where  $o(1)$  tends to 0 with  $\varepsilon$ . Hence

$$\int_V |\nabla^g \varphi_\varepsilon|^{\frac{2n}{n+1}} dv^g \leq C \left( \lambda_\varepsilon^{\frac{2n}{n+1}} \int_V |\varphi_\varepsilon|^{\frac{2n}{n-1}} dv^g + \varepsilon \int_V |\nabla^g \varphi_\varepsilon|^{\frac{2n}{n+1}} dv^g + o(1) \right). \quad (29)$$

This implies in particular that  $(\varphi_\varepsilon)$  is bounded in  $H_1^{\frac{2n}{n+1}}(V)$  and hence after passing to a subsequence  $(\varphi_\varepsilon)$  converges weakly to a limit  $\varphi$  in  $H_1^{\frac{2n}{n+1}}(V)$ .

The next step is to prove that  $\bar{\lambda} = \lim_{\varepsilon \rightarrow 0} \lambda_\varepsilon$  is not zero. To get a contradiction let us assume that  $\bar{\lambda} = 0$ . We then obtain from (29) that

$$\int_V |\nabla^g \varphi|^{\frac{2n}{n+1}} dv^g \leq \lim_{\varepsilon \rightarrow 0} \int_V |\nabla^g \varphi_\varepsilon|^{\frac{2n}{n+1}} dv^g = 0.$$

So  $\varphi$  is parallel and since  $D^g$  is invertible we conclude  $\varphi = 0$ , in other words  $(\varphi_\varepsilon)$  converges weakly to zero in  $H_1^{\frac{2n}{n+1}}(V)$ . As this space embeds compactly into  $L^{\frac{2n}{n+1}}(V)$  we have

$$\lim_{\varepsilon \rightarrow 0} \|\varphi_\varepsilon\|_{L^{\frac{2n}{n+1}}(V)} = \|\varphi\|_{L^{\frac{2n}{n+1}}(V)} = 0,$$

and hence  $(\varphi_\varepsilon)$  converges strongly to zero in  $H_1^{\frac{2n}{n+1}}(V)$ . As this space embeds continuously into  $L^{\frac{2n}{n-1}}(V)$  we conclude that the sequence converges strongly to zero in  $L^{\frac{2n}{n-1}}(V)$ . This is impossible since by Relation (24) we easily get that

$$\lim_{\varepsilon \rightarrow 0} \|\varphi_\varepsilon\|_{L^{\frac{2n}{n-1}}(V)} = 1.$$

From this contradiction we conclude

$$\bar{\lambda} > 0. \tag{30}$$

From (25) we have

$$\begin{aligned} \|D^g \varphi_\varepsilon\|_{L^{\frac{2n}{n+1}}(V)} &\leq \lambda_\varepsilon \|\varphi_\varepsilon\|_{L^{\frac{n-1}{n+1}}(V)}^{\frac{n-1}{n+1}} + \|A_{g_\varepsilon}^g(\nabla^g \varphi_\varepsilon)\|_{L^{\frac{2n}{n+1}}(V)} \\ &\quad + \|B_{g_\varepsilon}^g(\varphi_\varepsilon)\|_{L^{\frac{2n}{n+1}}(V)}. \end{aligned}$$

We already proved above that

$$\lim_{\varepsilon \rightarrow 0} \|B_{g_\varepsilon}^g(\varphi_\varepsilon)\|_{L^{\frac{2n}{n+1}}(V)} = 0.$$

Using Relation (27) we get similarly

$$\lim_{\varepsilon \rightarrow 0} \|A_{g_\varepsilon}^g(\nabla^g \varphi_\varepsilon)\|_{L^{\frac{2n}{n+1}}(V)} = 0.$$

Moreover since  $dv^{g_\varepsilon} = (1 + o(1)) dv^g$  it follows from (24) that

$$\lambda_\varepsilon \|\varphi_\varepsilon\|_{L^{\frac{n-1}{n+1}}(V)}^{\frac{n-1}{n+1}} = \lambda_\varepsilon (1 + o(1)).$$

We conclude


$$\|D^g \varphi_\varepsilon\|_{L^{\frac{2n}{n+1}}(V)} \leq \lambda_\varepsilon + o(1). \tag{31}$$

Starting from Equation (25) we can prove in a similar way that

$$\int_V \langle D^g \varphi_\varepsilon, \varphi_\varepsilon \rangle dv^g \geq \lambda_\varepsilon + o(1). \tag{32}$$

From (30), (31), and (32) it follows that  $\lambda \leq \lim_{\varepsilon \rightarrow 0} J(\varphi_\varepsilon) = \bar{\lambda}$ . This ends the demonstration of (22), which together with (19) proves Lemma 3.6.  $\square$



FIGURE 1. The function  $-\ln r \mapsto f(r)$ 

## 4. PROOFS

**4.1. Proof of Theorem 1.2.** This section is devoted to the proof of Theorem 1.2. Our goal is to construct a family of metrics  $(g_\theta)$  with  $0 < \theta < \theta_0$  which satisfies the conclusion of Theorem (1.2).

From Lemma 3.6 applied with  $V = M = M_1 \amalg M_2$  and  $S = W' = w_1(W \times \{0\}) \amalg w_2(W \times \{0\})$  we may assume that

$$g = h + dr^2 + r^2 \sigma^{n-k-1} \quad (33)$$

in a neighbourhood  $U(R_{\max})$  of  $W'$  where  $R_{\max} > 0$ . We fix numbers  $R_0, R_1 \in \mathbb{R}$  with  $R_{\max} > R_1 > R_0 > 0$  and we choose a function  $F : M \setminus W' \rightarrow \mathbb{R}^+$  such that

$$F(x) = \begin{cases} 1, & \text{if } x \in M_i \setminus U_i(R_1); \\ r_i^{-1} & \text{if } x \in U_i(R_0) \setminus W'. \end{cases}$$

We further choose  $\theta \in (0, R_0)$ , later we will let  $\theta \rightarrow 0$ . It is not difficult to see that there is a smooth function  $f : U(R_{\max}) \rightarrow \mathbb{R}$  (depending only on  $r$ ), real numbers  $\delta_1 = \delta_1(\theta)$  and  $\delta_2 = \delta_2(\theta)$  with  $\theta > \delta_2 > \delta_1 > 0$  and a real number  $A_\theta \in (\theta^{-1}, \delta_2^{-1})$  such that

$$f(x) = \begin{cases} -\ln r & \text{if } x \in U(R_{\max}) \setminus U(\theta); \\ \ln A_\theta & \text{if } x \in U(\delta_2), \end{cases}$$

and such that

$$\left| r \frac{df}{dr} \right| = \left| \frac{df}{d(\ln r)} \right| \leq 1,$$

and

$$\left\| r \frac{d}{dr} \left( r \frac{df}{dr} \right) \right\|_{L^\infty} = \left\| \frac{d^2 f}{d^2(\ln r)} \right\|_{L^\infty} \rightarrow 0$$

as  $\theta \rightarrow 0$ . It follows that  $\lim_{\theta \rightarrow 0} A_\theta = \infty$ .




FIGURE 2. Hierarchy of variables

After these choices we set  $\varepsilon := e^{-A_\theta \delta_1}$ . We assume that  $N$  is obtained from  $M$  by a connected sum along  $W$  with parameter  $\varepsilon$ , as explained in Section 2.2. In particular, recall that  $U_\varepsilon^N(s) = U(s) \setminus U(\varepsilon) / \sim$  for all  $s \geq \varepsilon$ . On the set  $U_\varepsilon^N(R_{\max}) = U(R_{\max}) \setminus U(\varepsilon) / \sim$  we define the variable  $t$  by

$$t := -\ln r_1 + \ln \varepsilon \leq 0$$

on  $U_1(R_{\max}) \setminus U(\varepsilon)$  and

$$t := \ln r_2 - \ln \varepsilon \geq 0$$

on  $U_2(R_{\max}) \setminus U(\varepsilon)$ . This implies

$$r_i = e^{|t| + \ln \varepsilon} = \varepsilon e^{|t|}.$$

The choices imply that  $t : U_\varepsilon^N(R_{\max}) \rightarrow \mathbb{R}$  is a smooth function with  $t \leq 0$  on  $U_\varepsilon^N(R_{\max}) \cap M_1$ ,  $t \geq 0$  on  $U_\varepsilon^N(R_{\max}) \cap M_2$ , and  $t = 0$  is the common boundary  $\partial U_1(\varepsilon)$  identified in  $N$  with  $\partial U_2(\varepsilon)$ . Then Equation (33) tells us that

$$r^{-2}g = \varepsilon^{-2}e^{-2|t|}h_i + dt^2 + \sigma^{n-k-1}.$$

Expressed in the new variable  $t$  we have

$$F(x) = \varepsilon^{-1}e^{-|t|}$$

if  $x \in U_\varepsilon^N(R_0) \setminus U_\varepsilon^N(\theta)$  or in other words if  $|t| + \ln \varepsilon \leq \ln R_0$ , and

$$f(t) = \begin{cases} -|t| - \ln \varepsilon & \text{if } |t| + \ln \varepsilon \in (\theta, R_{\max}), \\ \ln A_\theta & \text{if } |t| + \ln \varepsilon \leq \ln \delta_2, \end{cases}$$

and  $|df/dt| \leq 1$ ,  $\|d^2f/dt^2\|_{L^\infty} \rightarrow 0$ . After choosing a cut-off function  $\chi : \mathbb{R} \rightarrow [0, 1]$  such that  $\chi = 0$  on  $(-\infty, -1]$  and  $\chi = 1$  on  $[1, \infty)$ , we define

$$g_\theta(x) := \begin{cases} F^2 g_i & \text{if } x \in M_i \setminus U_i(\theta); \\ e^{2f(t)} h_i + dt^2 + \sigma^{n-k-1} & \text{if } x \in U_i(\theta) \setminus U_i(\delta_1); \\ A_\theta^2 \chi(A_\theta^{-1}t) h_2 + A_\theta^2 (1 - \chi(A_\theta^{-1}t)) h_1 + dt^2 + \sigma^{n-k-1} & \text{if } x \in U_i(\delta_1) \setminus U_i(\varepsilon). \end{cases}$$

(Recall that the  $h_i$  are defined as the pullback via  $w_i$  of the metric  $g_i$  on  $M_i$ , composed with restriction to  $W = W \times \{0\}$ .)

On  $U_\varepsilon^N(R_0)$  we write  $g_\theta$  as

$$g_\theta = \alpha_t^2 \tilde{h}_t + dt^2 + \sigma^{n-k-1},$$

where the metric  $\tilde{h}_t$  is defined for  $t \in \mathbb{R}$  by

$$\tilde{h}_t := \chi(A_\theta^{-1}t) h_2 + (1 - \chi(A_\theta^{-1}t)) h_1, \quad (34)$$

and where

$$\alpha_t := e^{f(t)}. \quad (35)$$

The rest of the proof consists of showing that  $(g_\theta)$  is the desired family of metrics. We first choose a sequence  $(\theta_i)_{i \in \mathbb{N}}$  converging to 0 so that  $\lim_{i \rightarrow \infty} \lambda_{\theta_i}$  exists. To avoid complicated notation we write  $\theta \rightarrow 0$  for the sequence  $(\theta_i)_{i \in \mathbb{N}}$  converging to zero and we will pass successively to subsequences without changing notation. Similarly  $\lim_{\theta \rightarrow 0} h(\theta)$  should be read as  $\lim_{i \rightarrow \infty} h(\theta_i)$ . We set  $\lambda := \lambda_{\min}^+(M_1 \amalg M_2, g)$ ,  $\lambda_\theta := \lambda_{\min}^+(N, g_\theta)$ , and  $\bar{\lambda} := \lim_{\theta \rightarrow 0} \lambda_\theta$ . Let  $J := J^g$  and  $J_\theta := J^{g_\theta}$  be the functionals associated respectively to  $g$  and  $g_\theta$ .

The easier part of the argument is to show that

$$\bar{\lambda} \leq \lambda. \quad (36)$$

For this let  $\alpha > 0$  be a small number. We choose a smooth cut-off function  $\chi_\alpha : M_1 \amalg M_2 \rightarrow [0, 1]$  such that  $\chi_\alpha = 1$  on  $M_1 \amalg M_2 \setminus U(2\alpha)$ ,  $|d\chi_\alpha| \leq 2/\alpha$ , and  $\chi_\alpha = 0$  on  $U(\alpha)$ . Let  $\psi$  be a smooth non-zero spinor such that  $J(\psi) \leq \lambda + \delta$  where  $\delta$  is a small positive number. On the support of  $\chi_\alpha$  the metrics  $g$  and  $g_\alpha$  are conformal since  $g_\theta = F^2g$  and hence by Formula (10) we have

$$\lambda_\theta \leq J_\theta \left( \chi_\alpha \beta_{g_\theta}^g (F^{-\frac{n-1}{2}} \psi) \right) = J(\chi_\alpha \psi)$$

for  $\theta < \alpha$ . Proceeding exactly as in the first part of the proof of Lemma 3.6 we show that  $\lim_{\alpha \rightarrow 0} J(\chi_\alpha \psi) = J(\psi) \leq \lambda + \delta$ . From this Relation (36) follows.

Now we turn to the more difficult part of the proof, that

$$\bar{\lambda} \geq \min\{\lambda, \Lambda_{n,k}\}. \quad (37)$$

By Proposition 2.3 we can assume that  $\lambda_\theta < \lambda_{\min}^+(S^n, \sigma^n)$  for all  $\theta$ , otherwise Relation (37) is trivial. From Theorem 2.4 we know that there exists a spinor field  $\psi_\theta \in \Gamma(\Sigma^{g_\alpha} N)$  of class  $C^2$  such that

$$\int_N |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} = 1$$

and

$$D^{g_\theta} \psi_\theta = \lambda_\theta |\psi_\theta|^{\frac{2}{n-1}} \psi_\theta. \quad (38)$$

We let  $x_\theta$  in  $N$  be such that  $|\psi_\theta(x_\theta)| = m_\theta$  where  $m_\theta := \|\psi_\theta\|_{L^\infty(N)}$ .

The proof continues divided in cases.

**Case I.** *The sequence  $(m_\theta)$  is not bounded.*

After taking a subsequence, we can assume that  $\lim_{\theta \rightarrow 0} m_\theta = \infty$ . We consider two subcases.

**Subcase I.1.** *There exists  $a > 0$  such that  $x_\theta \in N \setminus U^N(a)$  for an infinite number of  $\theta$ .*

We recall that  $N \setminus U^N(a) = N_\varepsilon \setminus U_\varepsilon^N(a) = M_1 \amalg M_2 \setminus U(a)$ . By taking a subsequence we can assume that there exists  $\bar{x} \in M_1 \amalg M_2 \setminus U(a)$  such that  $\lim_{\theta \rightarrow 0} x_\theta = \bar{x}$ . We let  $g'_\theta := m_\theta^{\frac{4}{n-1}} g_\theta$ . In a neighbourhood  $U$  of  $\bar{x}$  the metric  $g_\theta = F^2g$  does not depend on  $\theta$ . We apply Lemma 3.3 with  $O = U$ ,  $\alpha = \theta$ ,  $p_\alpha = x_\theta$ ,  $p = \bar{x}$ ,  $\gamma_\alpha = g_\theta = F^2g$ , and  $b_\alpha = m_\theta^{\frac{2}{n-1}}$ . Let  $r > 0$ . For  $\theta$  small enough Lemma 3.3 gives us diffeomorphisms

$$\Theta_\theta : B^n(r) \rightarrow B^{g_\theta}(x_\theta, m_\theta^{-\frac{2}{n-1}} r)$$

such that the sequence of metrics  $(\Theta_\theta^*(g'_\theta))$  tends to the Euclidean metric  $\xi^n$  in  $C^1(B^n(r))$ . We let  $\psi'_\theta := m_\theta^{-1}\psi_\theta$ . By (6) we then have

$$D^{g'_\theta}\psi'_\theta = \lambda_\theta|\psi'_\theta|^{\frac{2}{n-1}}\psi'_\theta$$

on  $B^{g_\theta}(x_\theta, m_\theta^{-\frac{2}{n-1}}r)$  and

$$\begin{aligned} \int_{B^{g_\theta}(x_\theta, m_\theta^{-\frac{2}{n-1}}r)} |\psi'_\theta|^{\frac{2n}{n-1}} dv^{g'_\theta} &= \int_{B^{g_\theta}(x_\theta, m_\theta^{-\frac{2}{n-1}}r)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} \\ &\leq \int_N |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} \\ &= 1. \end{aligned}$$

Here we used the fact that  $dv^{g'_\theta} = m_\theta^{\frac{2n}{n-1}} dv^{g_\theta}$ . Since

$$\Theta_\theta : (B^n(r), \Theta_\theta^*(g'_\theta)) \rightarrow (B^{g_\theta}(x_\theta, m_\theta^{-\frac{2}{n-1}}r), g'_\theta)$$

is an isometry we can consider  $\psi'_\theta$  as a solution of

$$D^{\Theta_\theta^*(g'_\theta)}\psi'_\theta = \lambda_\theta|\psi'_\theta|^{\frac{2}{n-1}}\psi'_\theta$$

on  $B^n(r)$  with  $\int_{B^n(r)} |\psi'_\theta|^{\frac{2n}{n-1}} dv^{\Theta_\theta^*(g'_\theta)} \leq 1$ . Since  $\|\psi_\theta\|_{L^\infty(B^n(r))} = |\psi'_\theta(0)| = 1$  we can apply Lemma 3.4 with  $V = \mathbb{R}^n$ ,  $\alpha = \theta$ ,  $g_\alpha = \Theta_\theta^*(g'_\theta)$ , and  $\psi_\alpha = \psi'_\theta$  (we may apply this lemma since each compact set of  $\mathbb{R}^n$  is contained in some ball  $B^n(r)$ ). This shows that there exists a spinor  $\psi$  of class  $C^1$  on  $(\mathbb{R}^n, \xi^n)$  which satisfies

$$D^{\xi^n}\psi = \bar{\lambda}|\psi|^{\frac{2}{n-1}}\psi.$$

Furthermore by (17) we have

$$\int_{B^n(r)} |\psi|^{\frac{2n}{n-1}} dv^{\xi^n} = \lim_{\theta \rightarrow 0} \int_{B^{g_\theta}(x_\theta, m_\theta^{-\frac{2}{n-1}}r)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} \leq 1$$

for any  $r > 0$ . We conclude that  $\int_{\mathbb{R}^n} |\psi|^{\frac{2n}{n-1}} dv^{\xi^n} \leq 1$ . Since  $|\psi(0)| = 1$  we also see that  $\psi$  is not identically zero. As  $(\mathbb{R}^n, \xi^n)$  and  $(S^n \setminus \{\text{pt}\}, \sigma^n)$  are conformal we can write  $\sigma^n = \Phi^2\xi^n$  for a positive function  $\Phi$ . We define  $\varphi := \Phi^{-\frac{n-1}{2}}\beta_{\sigma^n}^{\xi^n}\psi$ . By Equation (6) it follows that  $\varphi \in L^{\frac{2n}{n-1}}(S^n)$  is a solution of

$$D^{\sigma^n}\varphi = \bar{\lambda}|\varphi|^{\frac{2}{n-1}}\varphi \tag{39}$$

on  $S^n \setminus \{\text{pt}\}$  of class  $C^1$ . By Corollary 3.2 we know that  $\varphi$  can be extended to a weak solution of (39) on all  $S^n$  and by standard regularity theorems it follows that  $\varphi \in C^1(S^n)$ . Let  $J^{\sigma^n}$  be the functional associated to  $(S^n, \sigma^n)$ . By Equation (39) we have

$$\lambda_{\min}^+(S^n, \sigma^n) \leq J^{\sigma^n}(\varphi) = \bar{\lambda}$$

where the inequality comes from Proposition 2.3. We have proved Relation (37) in this subcase.

**Subcase I.2.** For all  $a > 0$  it holds that  $x_\theta \notin M_1 \amalg M_2 \setminus U(a)$  for  $\theta$  sufficiently small.

This means that  $x_\theta$  belongs to  $U^N(a)$  if  $\theta$  is sufficiently small. This subset is diffeomorphic to  $W \times I \times S^{n-k-1}$  where  $I$  is an interval. Through this diffeomorphism  $x_\theta$  can be written as

$$x_\theta = (y_\theta, t_\theta, z_\theta)$$

where  $y_\theta \in W$ ,  $t_\theta \in (-\ln R_0 + \ln \varepsilon, -\ln \varepsilon + \ln R_0)$ , and  $z_\theta \in S^{n-k-1}$ . By taking a subsequence we can assume that  $y_\theta$ ,  $\frac{t_\theta}{A_\theta}$ , and  $z_\theta$  converge respectively to  $y \in W$ ,  $T \in [-\infty, +\infty]$ , and  $z \in S^{n-k-1}$ . We apply Lemma 3.3 with  $V = W$ ,  $\alpha = \theta$ ,  $p_\alpha = y_\theta$ ,  $p = y$ ,  $\gamma_\alpha = \tilde{h}_{t_\theta}$ ,  $\gamma_0 = \tilde{h}_T$  (we define  $\tilde{h}_{-\infty} := h_1$  and  $\tilde{h}_{+\infty} := h_2$ ), and  $b_\alpha = m_\theta^{\frac{2}{n-1}} \alpha_{t_\theta}$ . The lemma provides diffeomorphisms

$$\Theta_\theta^y : B^k(r) \rightarrow B^{\tilde{h}_{t_\theta}}(y_\theta, m_\theta^{-\frac{2}{n-1}} \alpha_{t_\theta}^{-1} r)$$

for  $r > 0$  such that  $(\Theta_\theta^y)^*(m_\theta^{\frac{4}{n-1}} \alpha_{t_\theta}^2 \tilde{h}_{t_\theta})$  tends to the Euclidean metric  $\xi^k$  on  $B^k(r)$  as  $\theta \rightarrow 0$ . Next we apply Lemma 3.3 with  $V = S^{n-k-1}$ ,  $\alpha = \theta$ ,  $p_\alpha = z_\theta$ ,  $\gamma_\alpha = \gamma_0 = \sigma^{n-k-1}$ , and  $b_\alpha = m_\theta^{\frac{2}{n-1}}$ . For  $r' > 0$  we get the existence of diffeomorphisms

$$\Theta_\theta^z : B^{n-k-1}(r') \rightarrow B^{\sigma^{n-k-1}}(z_\theta, m_\theta^{-\frac{2}{n-1}} r')$$

such that  $(\Theta_\theta^z)^*(m_\theta^{\frac{4}{n-1}} \sigma^{n-k-1})$  converges to  $\xi^{n-k-1}$  on  $B^{n-k-1}(r')$  as  $\theta \rightarrow 0$ . For  $r, r', r'' > 0$  we define

$$\begin{aligned} U_\theta(r, r', r'') &:= B^{\tilde{h}_{t_\theta}}(y_\theta, m_\theta^{-\frac{2}{n-1}} \alpha_{t_\theta}^{-1} r) \times [t_\theta - m_\theta^{-\frac{2}{n-1}} r'', t_\theta + m_\theta^{-\frac{2}{n-1}} r''] \\ &\quad \times B^{\sigma^{n-k-1}}(z_\theta, m_\theta^{-\frac{2}{n-1}} r') \end{aligned}$$

and

$$\begin{aligned} \Theta_\theta : B^k(r) \times [-r'', r''] \times B^{n-k-1}(r') &\rightarrow U_\theta(r, r', r'') \\ (y, s, z) &\mapsto (\Theta_\theta^y(y), t(s), \Theta_\theta^z(z)), \end{aligned}$$

where  $t(s) := t_\theta + m_\theta^{-\frac{2}{n-1}} s$ . By construction  $\Theta_\theta$  is a diffeomorphism. As is readily seen

$$\Theta_\theta^*(m_\theta^{\frac{4}{n-1}} g_\theta) = (\Theta_\theta^y)^*(m_\theta^{\frac{4}{n-1}} \alpha_{t_\theta}^2 \tilde{h}_{t_\theta}) + ds^2 + (\Theta_\theta^z)^*(m_\theta^{\frac{4}{n-1}} \sigma^{n-k-1}). \quad (40)$$

By construction of  $\alpha_t$  one can verify that

$$\lim_{\theta \rightarrow 0} \left\| \frac{\alpha_{t_\theta}}{\alpha_t} - 1 \right\|_{C^1([t_\theta - m_\theta^{-\frac{2}{n-1}} r'', t_\theta + m_\theta^{-\frac{2}{n-1}} r''])} = 0$$

for all  $R > 0$  since  $\frac{df}{dt}$  and  $\frac{d^2f}{dt^2}$  are uniformly bounded. Moreover it is clear that

$$\lim_{\theta \rightarrow 0} \left| \tilde{h}_t - \tilde{h}_{t_\theta} \right|_{C^1(B^{\tilde{h}_{t_\theta}}(y_\theta, m_\theta^{-\frac{2}{n-1}} \alpha_{t_\theta}^{-1} R))} = 0$$

uniformly in  $t \in [t_\theta - m_\theta^{-\frac{2}{n-1}} r'', t_\theta + m_\theta^{-\frac{2}{n-1}} r'']$ . As a consequence

$$\lim_{\theta \rightarrow 0} \left| (\Theta_\theta^y)^* \left( m_\theta^{\frac{4}{n-1}} \left( \alpha_t^2 \tilde{h}_t - \alpha_{t_\theta}^2 \tilde{h}_{t_\theta} \right) \right) \right|_{C^1(B^k(r))} = 0$$

uniformly in  $t$ . This implies that the sequence  $(\Theta_\theta^y)^*(m_\theta^{\frac{4}{n-1}} \alpha_{t_\theta}^2 \tilde{h}_{t_\theta})$  tends to the Euclidean metric  $\xi^k$  in  $C^1(B^k(r))$  uniformly in  $t$  as  $\theta \rightarrow 0$ . From (40) we know that the sequence  $(\Theta_\theta^z)^*(m_\theta^{\frac{4}{n-1}} \sigma^{n-k-1})$  tends to the Euclidean metric  $\xi^{n-k-1}$  on  $B^{n-k-1}(r')$  as  $\theta \rightarrow 0$ . Returning to (40) we obtain that the sequence  $\Theta_\theta^*(m_\theta^{\frac{4}{n-1}} g_\theta)$  tends to  $\xi^n = \xi^k + ds^2 + \xi^{n-k-1}$  on  $B^k(r) \times [-r'', r''] \times B^{n-k-1}(r')$ . As in Subcase I.1 we apply Lemma 3.4 to get a spinor  $\psi$  of class  $C^1$  on  $\mathbb{R}^n$  which satisfies

$$D^{\xi^n} \psi = \bar{\lambda} |\psi|^{\frac{2}{n-1}} \psi$$

with  $\int_{B^n(r)} |\psi|^{\frac{2n}{n-1}} dx \leq 1$  for all  $r \in \mathbb{R}^+$ . Lemma 3.4 tells us that  $|\psi(0)| = 1$  so  $\psi$  does not vanish identically. As in Subcase I.1 we conclude that

$$\lambda \leq \lambda_{\min}^+(S^n, \sigma^n) \leq \bar{\lambda}.$$

This ends the proof of Theorem 1.2 in Case I.

**Case II.** *There exists a constant  $C_1$  such that  $m_\theta \leq C_1$  for all  $\theta$ .*

Again we consider two subcases.

**Subcase II.1.** *Assume that*

$$\liminf_{\theta \rightarrow 0} \int_{N \setminus U^N(a)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} > 0 \quad (41)$$

for some number  $a > 0$ .

Let  $K$  a compact subset such that  $K \subset M_1 \amalg M_2 \setminus W'$ . Choose a small number  $b$  such that  $K \subset M_1 \amalg M_2 \setminus U(2b) = N \setminus U^N(2b)$ . Let  $\chi \in C^\infty(M_1 \amalg M_2)$ ,  $0 \leq \chi \leq 1$ , be a cut-off function equal to 1 on  $M_1 \amalg M_2 \setminus U(2b)$  and equal to 0 on  $U(b)$ . Set  $\psi'_\theta := F^{\frac{n-1}{2}} (\beta_{g_\theta}^g)^{-1} \psi_\theta$ . Since  $g_\theta = F^2 g$  on the support of  $\chi$  we have

$$D^g \psi'_\theta = \lambda_\theta |\psi'_\theta|^{\frac{2}{n-1}} \psi'_\theta$$

on this set. For  $r > 0$  we have

$$\begin{aligned} & \int_{M_1 \amalg M_2} |D^g(\chi \psi'_\theta)|^r dv^g \\ &= \int_{M_1 \amalg M_2} \left| \text{grad}^g \chi \cdot \psi'_\theta + \chi \lambda_\theta |\psi'_\theta|^{\frac{2}{n-1}} \psi'_\theta \right|^r dv^g \\ &\leq 2^r \left( \int_{M_1 \amalg M_2} |\text{grad}^g \chi|^r |\psi'_\theta|^r dv^g + \lambda_\theta^r \int_{M_1 \amalg M_2} \chi^r |\psi'_\theta|^{\frac{(n+1)r}{n-1}} dv^g \right) \\ &\leq C. \end{aligned}$$

since  $m_\theta \leq C_1$ . Together with Relation (8) we get that the sequence  $(\chi \psi'_\theta)$  is bounded in  $H_1^1(M_1 \amalg M_2)$  for all  $r > 0$ . Proceeding as in the proof of Lemma 3.4 we get a  $C^1$  spinor  $\psi_0$  defined on  $K$  such that a subsequence of  $(\psi'_\theta)$  converges to  $\psi_0$  in  $C^0(K)$  and which satisfies

$$D^g \psi_0 = \bar{\lambda} |\psi_0|^{\frac{2}{n-1}} \psi_0. \quad (42)$$

Furthermore the convergence in  $C^0$  implies that

$$\int_K |\psi_0|^{\frac{2n}{n-1}} dv^g \leq \liminf_{\theta \rightarrow 0} \int_K |\psi'_\theta|^{\frac{2n}{n-1}} dv^g = \liminf_{\theta \rightarrow 0} \int_K |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} \leq 1.$$

Repeating the same for a sequence of compact sets which exhausts  $M_1 \amalg M_2 \setminus W'$  and taking a diagonal subsequence we can extend  $\psi_0$  to  $M_1 \amalg M_2 \setminus W'$ . Since  $\psi_0 \in L^{\frac{2n}{n-1}}(M_1 \amalg M_2 \setminus W') = L^{\frac{2n}{n-1}}(M_1 \amalg M_2)$  we can use Theorem 3.2 to extend  $\psi_0$  to a weak solution of Equation (42) on  $M_1 \amalg M_2$ . Note here that since  $D^g$  is invertible we have  $\bar{\lambda} > 0$ . By standard regularity theorems we conclude that

$\psi_0 \in C^1(M_1 \amalg M_2)$ . By (41) we have

$$\begin{aligned} \int_{M_1 \amalg M_2 \setminus U(a)} |\psi_0|^{\frac{2n}{n-1}} dv^g &= \lim_{\theta \rightarrow 0} \int_{M_1 \amalg M_2 \setminus U(a)} |\psi'_\theta|^{\frac{2n}{n-1}} dv^g \\ &= \lim_{\theta \rightarrow 0} \int_{M_1 \amalg M_2 \setminus U(a)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} \\ &> 0, \end{aligned}$$

and we conclude that  $\psi_0$  does not vanish identically. Equation (42) then leads to

$$\lambda \leq J(\psi_0) = \bar{\lambda} \left( \int_{M_1 \amalg M_2} |\psi_0|^{\frac{2n}{n-1}} dv^g \right)^{\frac{n+1}{n}-1} \leq \bar{\lambda},$$

which proves Theorem 1.2 in this case.

**Subcase II.2.** *We have*

$$\liminf_{\theta \rightarrow 0} \int_{N \setminus U^N(a)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} = 0 \quad (43)$$

for all  $a > 0$ .

This case is the most difficult one and we proceed in several steps. The assumption here is that we have a sequence  $(\theta_i)$  which tends to zero as  $i \rightarrow \infty$  with the property that the integral above tends to zero for all  $a > 0$ . We will abuse notation and write  $\lim_{\theta \rightarrow 0}$  for what should be a limit as  $i \rightarrow \infty$  or a limit of a subsequence.

For positive  $a$  and  $\theta$  let

$$\gamma_\theta(a) := \frac{\int_{N \setminus U^N(a)} |\psi_\theta|^2 dv^{g_\theta}}{\int_{U^N(a)} |\psi_\theta|^2 dv^{g_\theta}}$$

The first step is to establish an estimate for  $\gamma_\theta(a)$ .

**Step 1.** *There is a constant  $C_0$  so that*

$$1 \leq C_0 \left( \gamma_\theta(a) + \|\psi_\theta\|_{L^\infty(U^N(2a))}^{\frac{4}{n-1}} \right) \quad (44)$$

for all  $a > 0$ .

Let  $\chi \in C^\infty(N)$ ,  $0 \leq \chi \leq 1$ , be a cut-off function with  $\chi = 1$  on  $U^N(a)$  and  $\chi = 0$  on  $N \setminus U^N(2a) = M_1 \amalg M_2 \setminus U(2a)$ . Since the definitions of  $U^N(a)$  and  $U(a)$  use the distance to  $W'$  for the metric  $g$  we can and do assume that  $|d\chi|_g \leq 2/a$ . For the metric  $g_\theta$  this gives

$$|d\chi|_{g_\theta} = F^{-1}|d\chi|_g = r|d\chi|_g \leq 2a \frac{2}{a} = 4.$$

From Lemma 3.5 and Equation (38) it follows that

$$\begin{aligned}
\frac{(n-k-1)^2}{4} &\leq \frac{\int_N |D^{g_\theta}(\chi\psi_\theta)|^2 dv^{g_\theta}}{\int_N |\chi\psi_\theta|^2 dv^{g_\theta}} \\
&= \frac{\int_N |d\chi|_{g_\theta}^2 |\psi_\theta|^2 dv^{g_\theta} + \lambda_\theta^2 \int_N \chi^2 |\psi_\theta|^{\frac{2(n+1)}{n-1}} dv^{g_\theta}}{\int_N |\chi\psi_\theta|^2 dv^{g_\theta}} \\
&\leq \frac{16 \int_{U^N(2a) \setminus U^N(a)} |\psi_\theta|^2 dv^{g_\theta} + \lambda_\theta^2 \|\psi_\theta\|_{L^\infty(U^N(2a))}^{\frac{4}{n-1}} \int_N |\chi\psi_\theta|^2 dv^{g_\theta}}{\int_N |\chi\psi_\theta|^2 dv^{g_\theta}} \\
&\leq \frac{16 \int_{U^N(2a) \setminus U^N(a)} |\psi_\theta|^2 dv^{g_\theta}}{\int_{U^N(a)} |\psi_\theta|^2 dv^{g_\theta}} + \lambda_\theta^2 \|\psi_\theta\|_{L^\infty(U^N(2a))}^{\frac{4}{n-1}} \\
&\leq 16\gamma_\theta(a) + \lambda_\theta^2 \|\psi_\theta\|_{L^\infty(U^N(2a))}^{\frac{4}{n-1}}.
\end{aligned}$$

Using that  $\lambda_\theta \leq \lambda_{\min}^+(S^n, \sigma^n)$  by Proposition 2.3 we obtain Relation (44) with

$$C_0 := \frac{4}{(n-k-1)^2} \max \{16, \lambda_{\min}^+(S^n)^2\}.$$

This ends the proof of Step 1.

**Step 2.** *There exist a sequence of positive numbers  $(a_\theta)$  which tends to 0 with  $\theta$  and constants  $0 < m < M$  such that*

$$m \leq \|\psi_\theta\|_{L^\infty(U^N(2a_\theta))} \leq M \quad (45)$$

for all  $\theta$ .

By (43) we have

$$\lim_{\theta \rightarrow 0} \int_{N \setminus U^N(a)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} = 0$$

for all  $a > 0$ . Since  $\text{Vol}(N \setminus U^N(a), g_\theta)$  does not depend on  $\theta$  if  $\theta < a$  it follows that

$$\lim_{\theta \rightarrow 0} \left( \int_{N \setminus U^N(a)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} \right)^{\frac{n-1}{n}} \text{Vol}(N \setminus U^N(a), g_\theta)^{\frac{1}{n}} = 0$$

for all  $a$ . Hence we can take a sequence  $(a_\theta)$  which tends sufficiently slowly to 0 so that

$$\lim_{\theta \rightarrow 0} \left( \int_{N \setminus U^N(a_\theta)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} \right)^{\frac{n-1}{n}} \text{Vol}(N \setminus U^N(a_\theta), g_\theta)^{\frac{1}{n}} = 0. \quad (46)$$

Using the Hölder inequality we get

$$\begin{aligned}
\gamma_\theta(a_\theta) &= \frac{\int_{N \setminus U^N(a_\theta)} |\psi_\theta|^2 dv^{g_\theta}}{\int_{U^N(a_\theta)} |\psi_\theta|^2 dv^{g_\theta}} \\
&\leq \frac{\left( \int_{N \setminus U^N(a_\theta)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} \right)^{\frac{n-1}{n}} \text{Vol}(N \setminus U^N(a_\theta), g_\theta)^{\frac{1}{n}}}{\|\psi_\theta\|_{L^\infty(U^N(a_\theta))}^{-\frac{2}{n-1}} \int_{U^N(a_\theta)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta}}.
\end{aligned}$$



The numerator of this expression tends to 0 by Relation (46). Further by (46) we have

$$\begin{aligned} \lim_{\theta \rightarrow 0} \int_{U^N(a_\theta)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} &= \lim_{\theta \rightarrow 0} \int_N |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} - \int_{N \setminus U^N(a_\theta)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} \\ &= 1. \end{aligned}$$

Together with the fact that  $\|\psi_\theta\|_{L^\infty(U^N(a_\theta))} \leq m_\theta \leq C_1$  we obtain that

$$\lim_{\theta \rightarrow 0} \gamma_\theta(a_\theta) = 0.$$

From Relation (44) applied with  $a = a_\theta$  we know that  $\|\psi_\theta\|_{L^\infty(U^N(2a_\theta))}$  is bounded from below. Moreover, by the assumption of Case II we have that  $\|\psi_\theta\|_{L^\infty(U^N(2a_\theta))} \leq m_\theta \leq C_1$ . This finishes the proof of Step 2.

**Step 3.** *We have*

$$\bar{\lambda} \geq \Lambda_{n,k}.$$

Let  $x_\theta$  be a point in the closure of  $U^N(2a_\theta)$  such that  $|\psi_\theta(x_\theta)| = \|\psi_\theta\|_{L^\infty(U^N(2a_\theta))}$ . As in Subcase I.2 we write  $x_\theta = (y_\theta, t_\theta, z_\theta)$  where  $y_\theta \in W$ ,  $t_\theta \in (-\ln R_0 + \ln \varepsilon, -\ln \varepsilon + \ln R_0)$ , and  $z_\theta \in S^{n-k-1}$ . By restricting to a subsequence we can assume that  $y_\theta$ ,  $t_\theta/A_\theta$ , and  $z_\theta$  converge respectively to  $y \in W$ ,  $T \in [-\infty, +\infty]$ , and  $z \in S^{n-k-1}$ . We apply Lemma 3.3 with  $V = W$ ,  $\alpha = \theta$ ,  $p_\alpha = y_\theta$ ,  $p = y$ ,  $\gamma_\alpha = \tilde{h}_{t_\theta}$ ,  $\gamma_0 = \tilde{h}_T$ , and  $b_\alpha = \alpha_{t_\theta}$  (recall that  $\tilde{h}_t$  and  $\alpha_t$  were defined in (34) and (35)) and conclude that there is a diffeomorphism

$$\Theta_\theta^y : B^k(r) \rightarrow B^{\tilde{h}_{t_\theta}}(y_\theta, \alpha_{t_\theta}^{-1}r)$$

for  $r > 0$  such that  $(\Theta_\theta^y)^*(\alpha_{t_\theta}^2 \tilde{h}_{t_\theta})$  converges to the Euclidean metric  $\xi^k$  on  $B^k(r)$ . For  $r, r' > 0$  we define

$$U_\theta(r, r') := B^{\tilde{h}_{t_\theta}}(y_\theta, \alpha_{t_\theta}^{-1}r) \times [t_\theta - r', t_\theta + r'] \times S^{n-k-1}$$

and

$$\begin{aligned} \Theta_\theta : B^k(r) \times [-r', r'] \times S^{n-k-1} &\rightarrow U_\theta(r, r') \\ (y, s, z) &\mapsto (\Theta_\theta^y(y), t(s), z), \end{aligned}$$

where  $t(s) := t_\theta + s$ . By construction  $\Theta_\theta$  is a diffeomorphism. Since  $g_\theta = \alpha_{t_\theta}^2 \tilde{h}_{t_\theta} + dt^2 + \sigma^{n-k-1}$  we see that

$$\Theta_\theta^*(g_\theta) = \frac{\alpha_{t_\theta}^2}{\alpha_{t_\theta}^2} (\Theta_\theta^y)^*(\alpha_{t_\theta}^2 \tilde{h}_{t_\theta}) + ds^2 + \sigma^{n-k-1}. \quad (47)$$

We will now find the limit of  $\Theta_\theta^*(g_\theta)$  in the  $C^1$  topology. We define  $c := \lim_{\theta \rightarrow 0} f'(t_\theta)$ .

**Lemma 4.1.** *The sequence of metrics  $\Theta_\theta^*(g_\theta)$  tends to*

$$G_c = \eta_c^{k+1} + \sigma^{n-k-1} = e^{2cs} \xi^k + ds^2 + \sigma^{n-k-1}$$

*in  $C^1$  on  $B^k(r) \times [-r', r'] \times S^{n-k-1}$  for fixed  $r, r' > 0$ .*

*Proof.* Recall that  $\alpha_t = e^{f(t)}$ . The intermediate value theorem tells us that

$$|f(t) - f(t_\theta) - f'(t_\theta)(t - t_\theta)| \leq \frac{r'^2}{2} \max_{\xi \in [t_\theta - r', t_\theta + r']} |f''(\xi)|$$

for all  $t \in [t_\theta - r', t_\theta + r']$ . On the other hand we assume that  $f''(t) \rightarrow 0$  as  $\theta \rightarrow 0$ , so

$$\|f(t) - f(t_\theta) - f'(t_\theta)(t - t_\theta)\|_{C^0([t_\theta - r', t_\theta + r'])} \rightarrow 0$$

as  $\theta \rightarrow 0$  (and  $r'$  fixed). Furthermore

$$\begin{aligned} \left| \frac{d}{dt} (f(t) - f(t_\theta) - f'(t_\theta)(t - t_\theta)) \right| &= |f'(t) - f'(t_\theta)| \\ &= \left| \int_{t_\theta}^t f''(s) ds \right| \\ &\leq r' \max_{\xi \in [t_\theta - r', t_\theta + r']} |f''(\xi)| \\ &\rightarrow 0 \end{aligned}$$

as  $\theta \rightarrow 0$ . Together with  $c = \lim_{\theta \rightarrow 0} f'(t_\theta)$  we have

$$\|f(t) - f(t_\theta) - c(t - t_\theta)\|_{C^1([t_\theta - r', t_\theta + r'])} \rightarrow 0.$$

Exponentiation of functions is a continuous map

$$C^1([t_\theta - r', t_\theta + r']) \ni \tilde{f} \mapsto \exp \circ \tilde{f} \in C^1([t_\theta - r', t_\theta + r']).$$

Hence

$$\left\| \frac{\alpha_t}{\alpha_{t_\theta}} - e^{c(t-t_\theta)} \right\|_{C^1([t_\theta - r', t_\theta + r'])} = \left\| e^{f(t)-f(t_\theta)} - e^{c(t-t_\theta)} \right\|_{C^1([t_\theta - r', t_\theta + r'])} \rightarrow 0$$

as  $\theta \rightarrow 0$ . We now write  $\alpha_t^2 \tilde{h}_t = \alpha_t^2 (\tilde{h}_t - \tilde{h}_{t_\theta}) + \frac{\alpha_t^2}{\alpha_{t_\theta}^2} \alpha_{t_\theta}^2 \tilde{h}_{t_\theta}$ . Using the fact that

$$\lim_{\theta \rightarrow 0} \left\| \tilde{h}_t - \tilde{h}_{t_\theta} \right\|_{C^1(B_{\tilde{h}_{t_\theta}}(y_\theta, \alpha_{t_\theta}^{-1} R))} = 0$$

uniformly for  $t \in [t_\theta - r', t_\theta + r']$  we get that the sequence  $\frac{\alpha_t^2}{\alpha_{t_\theta}^2} (\Theta_\theta^y)^* (\alpha_{t_\theta}^2 \tilde{h}_t)$  tends to  $e^{2cs} \xi^k$  in  $C^1$  on  $B^k(r)$ . Going back to Relation (47) this proves Lemma 4.1.  $\square$

We continue with the proof of Step 3. As in subcases I.1 and I.2 we apply Lemma 3.4 with  $(V, g) = (\mathbb{R}^{k+1} \times S^{n-k-1}, G_c)$ ,  $\alpha = \theta$ , and  $g_\alpha = \Theta_\theta^*(g_\theta)$  (we can apply this lemma since any compact subset of  $\mathbb{R}^{k+1} \times S^{n-k-1}$  is contained in some  $B^k(r) \times [-r', r'] \times S^{n-k-1}$ ). We obtain a  $C^1$  spinor  $\psi$  which is a solution of

$$D^{G_c} \psi = \bar{\lambda} |\psi|^{\frac{2}{n-1}} \psi$$

on  $(\mathbb{R}^{k+1} \times S^{n-k-1}, G_c)$ . From (17) it follows that

$$\int_{\mathbb{R}^{k+1} \times S^{n-k-1}} |\psi|^{\frac{2n}{n-1}} dv^{G_c} \leq 1.$$

From (16) it follows that  $\psi \in L^\infty(\mathbb{R}^{k+1} \times S^{n-k-1})$ , and from (16) and (45) it follows that  $\psi$  does not vanish identically. We want to show that  $\psi \in L^2(\mathbb{R}^{k+1} \times S^{n-k-1})$ . From (17) we get that

$$\begin{aligned} \int_{B^k(r) \times [-r', r'] \times S^{n-k-1}} |\psi|^2 dv^{G_c} &= \lim_{\theta \rightarrow 0} \int_{U_\theta(r, r')} |\psi_\theta|^2 dv^{g_\theta} \\ &\leq \lim_{\theta \rightarrow 0} \int_{U^N(a)} |\psi_\theta|^2 dv^{g_\theta} \end{aligned} \tag{48}$$

for some fixed number  $a > 0$  independent of  $r$ ,  $r'$  and  $\theta$ . Let  $\chi$  be defined as in Step 1. Using the Hölder inequality, Lemma 3.5, and Equation (38) we see that

$$\begin{aligned} & \frac{(n-k-1)^2}{4} \\ & \leq \frac{\int_N |D^{g_\theta}(\chi\psi_\theta)|^2 dv^{g_\theta}}{\int_N |\chi\psi_\theta|^2 dv^{g_\theta}} \\ & = \frac{\int_N |d\chi|_{g_\theta}^2 |\psi_\theta|^2 dv^{g_\theta} + \lambda_\theta^2 \int_N \chi^2 |\psi_\theta|^{\frac{2(n+1)}{n-1}} dv^{g_\theta}}{\int_N |\chi\psi_\theta|^2 dv^{g_\theta}} \\ & \leq \frac{16 \int_{U^N(2a) \setminus U^N(a)} |\psi_\theta|^2 dv^{g_\theta} + \lambda_\theta^2 \|\psi_\theta\|_{L^\infty(U^N(2a))}^{\frac{2}{n-1}} \int_{U^N(2a)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta}}{\int_{U^N(a)} |\psi_\theta|^2 dv^{g_\theta}}. \end{aligned}$$

We have

$$\lambda_\theta^2 \|\psi_\theta\|_{L^\infty(U^N(2a))}^{\frac{2}{n-1}} \int_{U^N(2a)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} \leq \lambda_{\min}^+(S^n, \sigma^n)^2 C_1^{\frac{2}{n-1}}$$

and

$$\begin{aligned} & \int_{U^N(2a) \setminus U^N(a)} |\psi_\theta|^2 dv^{g_\theta} \\ & \leq \left( \int_{U^N(2a) \setminus U^N(a)} |\psi_\theta|^{\frac{2n}{n-1}} dv^{g_\theta} \right)^{\frac{n-1}{n}} \text{Vol}(U^N(2a) \setminus U^N(a), g_\theta)^{\frac{1}{n}} \\ & \leq \text{Vol}(U^N(2a) \setminus U^N(a), g_\theta)^{\frac{1}{n}}. \end{aligned}$$

Since  $g_\theta$  does not depend on  $\theta$  on  $U^N(2a) \setminus U^N(a)$  for  $\theta < a$ , we get the existence of a constant  $C$  such that

$$\frac{(n-k-1)^2}{4} \leq \frac{C}{\int_{U^N(a)} |\psi_\theta|^2 dv^{g_\theta}}.$$

Together with (48) we obtain that

$$\int_{B^k(r) \times [-r', r'] \times S^{n-k-1}} |\psi|^2 dv^{G_c} \leq C$$

where  $C$  is independent of  $r$  and  $r'$ . This proves that  $\psi \in L^2(\mathbb{R}^{k+1} \times S^{n-k-1})$ . Since the spinor  $\psi$  is non-zero and

$$\psi \in L^\infty(\mathbb{R}^{k+1} \times S^{n-k-1}) \cap C_{\text{loc}}^1(\mathbb{R}^{k+1} \times S^{n-k-1}) \cap L^2(\mathbb{R}^{k+1} \times S^{n-k-1})$$

with

$$\int_{\mathbb{R}^{k+1} \times S^{n-k-1}} |\psi|^{\frac{2n}{n-1}} dv^{G_c} \leq 1$$

we get that  $\bar{\lambda} \geq \Lambda_{n,k}$  by the definition of  $\Lambda_{n,k}$ . This ends the proof of this subcase and the proof of Theorem 1.2.

**4.2. Proof of Theorem 1.1.** We prove Theorem 1.1 by contradiction. Assume that there is a sequence  $c_i \in [-1, 1]$ ,  $i \in \mathbb{N}$ , for which

$$\lim_{i \rightarrow \infty} \widetilde{\lambda}_{\min}^+(\mathbb{R}^{k+1} \times S^{n-k-1}, G_{c_i}) = 0.$$

After removing the indices  $i$  for which  $\widetilde{\lambda}_{\min}^+$  is infinite we have for all  $i$  a solution of

$$D^{G_{c_i}} \psi_i = \lambda_i |\psi_i|^{\frac{2}{n-1}} \psi_i \quad (49)$$

where  $\lambda_i \rightarrow 0$  as  $i \rightarrow \infty$ . Moreover, the spinors  $\psi_i$  are in  $L^\infty \cap L^2 \cap C_{\text{loc}}^1$  and

$$\int_{\mathbb{R}^{k+1} \times S^{n-k-1}} |\psi_i|^{\frac{2n}{n-1}} dv^{G_{c_i}} \leq 1.$$

Let  $m_i := \|\psi_i\|_{L^\infty}$ . We cannot assume that  $m_i$  is attained, but since  $(\mathbb{R}^{k+1} \times S^{n-k-1}, G_{c_i})$  is a symmetric space we can compose  $\psi_i$  with isometries so that  $|\psi_i(P)| > m_i/2$  for some fixed point  $P \in \mathbb{R}^{k+1} \times S^{n-k-1}$ . First we prove that

$$\lim_{i \rightarrow \infty} m_i = \infty. \quad (50)$$

By Lemma 3.5 and Equation (49) we have

$$\begin{aligned} \frac{(n-k-1)^2}{4} &\leq \frac{\int_{\mathbb{R}^{k+1} \times S^{n-k-1}} |D^{G_{c_i}} \psi_i|^2 dv^{G_{c_i}}}{\int_{\mathbb{R}^{k+1} \times S^{n-k-1}} |\psi_i|^2 dv^{G_{c_i}}} \\ &\leq \frac{\lambda_i^2 \int_{\mathbb{R}^{k+1} \times S^{n-k-1}} |\psi_i|^{\frac{2(n+1)}{n-1}} dv^{G_{c_i}}}{\int_{\mathbb{R}^{k+1} \times S^{n-k-1}} |\psi_i|^2 dv^{G_{c_i}}} \\ &\leq \lambda_i^2 m_i^{\frac{4}{n-1}}. \end{aligned}$$

Since  $\lambda_i$  tends to zero this proves (50). Restricting to subsequence we can assume that  $\lim_{i \rightarrow \infty} c_i$  exists and we denote this limit by  $c \in [-1, 1]$ . We apply Lemma 3.3 with  $\alpha = 1/i$ ,  $(V, \gamma_\alpha) = (\mathbb{R}^{k+1} \times S^{n-k-1}, G_{c_i})$ ,  $(V, \gamma_0) = (\mathbb{R}^{k+1} \times S^{n-k-1}, G_c)$ ,  $p_\alpha = p = P$ , and  $b_\alpha = m_i^{\frac{2}{n-1}}$ . For  $r > 0$  we obtain a diffeomorphism

$$\Theta_i : B^n(r) \rightarrow B^{G_{c_i}}(P, m_i^{\frac{2}{n-1}} r)$$

such that  $\Theta_i^*(m_i^{\frac{4}{n-1}}(G_{c_i}))$  tends to the Euclidean metric  $\xi^n$  on  $B^n(r)$ . Proceeding as in Subcase I.1 of Theorem 1.2 we construct a non-zero spinor  $\psi$  belonging to  $L^{\frac{2n}{n-1}}(\mathbb{R}^n)$  such that

$$D^{\xi^n} \psi = \lim_{i \rightarrow \infty} \lambda_i |\psi|^{\frac{2}{n-1}} \psi = 0.$$

Again as in Subcase I.1 of Theorem 1.2 we get  $0 \geq \lambda_{\min}^+(S^n, \sigma^n)$ , which is false. This proves Theorem 1.1.

## REFERENCES

- [1] B. Ammann, *The smallest Dirac eigenvalue in a spin-conformal class and cmc-immersions*, Preprint, 2003.
- [2] ———, *A spin-conformal lower bound of the first positive Dirac eigenvalue*, Diff. Geom. Appl. **18** (2003), 21–32.
- [3] ———, *A variational problem in conformal spin geometry*, Habilitationsschrift, Universität Hamburg, 2003.
- [4] B. Ammann, M. Dahl, and E. Humbert, *Surgery and harmonic spinors*, Preprint, ArXiv math.DG/0606224, 2006, to appear in Adv. Math. **220** (2009) no. 2, 523–539.
- [5] ———, *Smooth Yamabe invariant and surgery*, Preprint, ArXiv 0804.1418, 2008.

- [6] B. Ammann, J. F. Grosjean, E. Humbert, and B. Morel, *A spinorial analogue of Aubin's inequality*, Math. Z. **260** (2008), 127–151.
- [7] B. Ammann and E. Humbert, *The spinorial  $\tau$ -invariant and 0-dimensional surgery*, J. reine angew. Math., **624** (2008), 27–50.
- [8] H. Baum, *Spin-Strukturen und Dirac-Operatoren über pseudoriemannschen Mannigfaltigkeiten*, Teubner Verlag, 1981.
- [9] J.-P. Bourguignon and P. Gauduchon, *Spineurs, opérateurs de Dirac et variations de métriques*, Comm. Math. Phys. **144** (1992), 581–599.
- [10] T. Friedrich, *Dirac Operators in Riemannian Geometry*, Graduate Studies in Mathematics 25, AMS, Providence, Rhode Island, 2000.
- [11] D. Gilbarg and N. Trudinger, *Elliptic partial differential equations of second order*, Grundlehren der mathematischen Wissenschaften, no. 224, Springer-Verlag, 1977.
- [12] O. Hijazi, *A conformal lower bound for the smallest eigenvalue of the Dirac operator and Killing spinors*, Comm. Math. Phys. **104** (1986), 151–162.
- [13] ———, *Première valeur propre de l'opérateur de Dirac et nombre de Yamabe*, C. R. Acad. Sci. Paris t. **313**, Série I (1991), 865–868.
- [14] ———, *Spectral properties of the Dirac operator and geometrical structures.*, Ocampo, Hernan (ed.) et al., Geometric methods for quantum field theory. Proceedings of the summer school, Villa de Leyva, Colombia, July 12-30, 1999. Singapore: World Scientific. 116–169 , 2001.
- [15] N. Hitchin, *Harmonic spinors*, Adv. Math. **14** (1974), 1–55.
- [16] R. C. Kirby, *The topology of 4-manifolds*, Lecture Notes in Mathematics, vol. 1374, Springer-Verlag, Berlin, 1989.
- [17] O. Kobayashi, *Scalar curvature of a metric with unit volume*, Math. Ann. **279** (1987), no. 2, 253–265.
- [18] A. A. Kosinski, *Differential manifolds*, Pure and Applied Mathematics, vol. 138, Academic Press Inc., Boston, MA, 1993.
- [19] H. B. Lawson and M.-L. Michelsohn, *Spin geometry*, Princeton University Press, Princeton, 1989.
- [20] J. Lott, *Eigenvalue bounds for the Dirac operator*, Pacific J. of Math. **125** (1986), 117–126.
- [21] R. Schoen, *Variational theory for the total scalar curvature functional for Riemannian metrics and related topics*, Topics in calculus of variations, Lect. 2nd Sess., Montecatini/Italy 1987, Lect. Notes Math. 1365, 120-154, 1989.
- [22] R. E. Stong, *Notes on cobordism theory*, Mathematical notes, Princeton University Press, Princeton, N.J., 1968.

NWF I – MATHEMATIK, UNIVERSITÄT REGENSBURG, 93040 REGENSBURG, GERMANY  
*E-mail address:* `bernd.ammann@mathematik.uni-regensburg.de`

INSTITUTIONEN FÖR MATEMATIK, KUNGLIGA TEKNISKA HÖGSKOLAN, 100 44 STOCKHOLM, SWEDEN  
*E-mail address:* `dahl@math.kth.se`

INSTITUT ÉLIE CARTAN, BP 239, UNIVERSITÉ DE NANCY 1, 54506 VANDOEUVRE-LÈS-NANCY CEDEX, FRANCE  
*E-mail address:* `humbert@iecn.u-nancy.fr`